

[색상에 유의하여 인쇄 바랍니다]

디자인 사양	자재명	코스판주사20mg_설명서(수)-싱가폴
주의사항 1. JPG 파일(그림파일)이나 PDF파일을 첨부하오니 그림유실, 데이터 오류 등 체크해 주시기 바랍니다. 2. 감리 일자는 일주일 전 협의바랍니다. 3. 감리를 가지 않을 경우 사전에 담당자와 협의하여 사진을 통해 인쇄 상태를 확인 할 수 있어야 합니다. 4. 파일에 문제가 있을 시 임의 변경 하지 마시고 연락 주시기 바랍니다. 5. 인쇄도수에 'P'를 표기한 경우 'PANTONE'을 의미합니다.	최종수정일	2024.03.13
	규격(장폭고)mm	110*245mm
	재질	모조지 70g
	양면여부	양면인쇄
	비고	-
위 사항을 지키지 않아 발생하는 문제는 본사에서 책임지지 않습니다.	문의전화	삼진제약 커뮤니케이션팀 02-3140-0693
인쇄도수		
1도		
	BLACK	

Antispasmodic agent

Cospan INJECTION 20Mmg/ml (1ml)

(Hyoscine –N–Butylbromide 20mg)

■ Composition

One ampoule (1ml) contains 20mg of Hyoscine-N-Butylbromide as active substance with water for injection.

■ Description

Colorless and transparent ampoule containing colorless and clear solution

■ Properties

Cospan exerts antispasmodic effect on smooth muscles of gastro-intestinal, biliary and urinary tract by acting selectively on cholinergic nerves in viscera paries.

■ Pharmacokinetics

Hyoscine is readily absorbed from the gastro intestinal tract following oral doses of the hydrobromide. It is almost entirely metabolised, probably in the liver; only a small proportion of an oral dose has been reported to be excreted unchanged in the urine. It crosses the blood-brain barrier and has been stated to cross the placenta. Hyoscine is also well absorbed following application to the skin.

The quaternary derivatives, such as the butyibromide or methobromide are poorly absorbed from the gastro-intestinal tract and do not readily pass the blood brain barrier.

■ Pharmacodynamics

Cospan Injection is an antispasmodic agent which relaxes smooth muscle of the organs of the abdominal and pelvic cavities. It is believed to act predominantly on the intramural parasympathetic ganglia of these organs.

■ Indications

Gastro-duodenal ulcer, cardiospasm, pylorospasm, esophagospasm, gastritis, duodenitis, enteritis intestinal colic, spastic constipation, cholecystitis, cholangitis, cholelithiasis, syndromes following gastectomy or cholecystectomy, dyskinesia of bile duct, lithangiuria, cystitis, functional diarrhea, dysmenorrhea, lower uterine spasms during delivery, urinary bladder spasms caused by apparatus insertion. Premanagement of gastro-intestinal radiography and endoscopy.

■ Recommended Dosage

Once 10mg to 20mg I.V or I.M daily.

■ Route of Administration

Intravenous (IV) + Intramuscular (IM)

■ Contraindications

Glaucoma, prostatomegaly, pyloric stenosis and paralytic lieus. Cospan Injection should not be administered to patients with tachycardia.

■ Warning and Precautions

General Precaution:

- 1) In patients with severe cardiac diseases, coronary arterial insufficiency or cardiac insufficiency, it should be administered following physician's direction.
- 2) It should be administered carefully when driving cars or operating engines.

Warnings and Precautions

Cospan Injection can cause tachycardia, hypotension and anaphylaxis, therefore use with caution in patients with cardiac conditions such as cardiac failure, coronary heart disease or cardiac arrhythmia and patients with cardiac cardiovascular disease (e.g. acute myocardial infarction, hypertension and conditions associated with tachycardia or hypertension), and in cardiac surgery).

Monitoring of these patients is advised. Emergency equipment and personnel trained in its use must be readily available.

■ Usage during Pregnancy and Lactation

It should only be used if the potential benefit justifies the potential risk to the fetus.

■ Adverse Effects / Undesirable Effects

Side Effects

Constipation, thirst, dysuria, mydriasis, cycloplegia, flush, vertigo, vomiting anxiety, agitation hallucination, orthostatic hypotension, eruption, tachycardia, hyperthermia.

Adverse Effects/Undesirable Effects:

- Immune system disorders

Not known: anaphylactic shock including cases with fatal outcome, anaphylactic reactions.

- Cardiac disorders

Common: tachycardia

■ Overdose and Treatment

Symptoms

Serious signs of poisoning following acute overdosage have not been observed in man. In the case of overdosage, anticholinergic symptoms such as urinary retention, dry mouth, reddening of the skin, tachycardia, inhibition of gastrointestinal motility and transient visual disturbances may occur, and Cheynes-Stokes respiration has been reported.

Therapy

Symptoms of Cospan Injection overdosage respond to parasympathomimetics. For patients with glaucoma, pilocarpine should be given locally. Cardiovascular complications should be treated according to usual therapeutic principles. In case of respiratory paralysis: intubation, artificial respiration should be considered. Catheterisation may be required for urinary retention.

In addition, appropriate supportive measures should be used as required.

■ Incompatibilities

The effects of antimuscarinic agents may be enhanced by the consistent administration of other drugs with antimuscarinic properties, such as amantadine, some antihistamines, butyrophenones and phenothiazine and tricyclic antidepressants. The reduction in gastric motility caused by antimuscarinic agents may affect the absorption of other drugs.

■ Storage Conditions

Store in light-resistant, hermetic container below 30°C.

■ Dosage forms and Packaging available

1ml(20mg) X 6 Ampoules

■ Product Registration Holder

The Zyfas Medical Co.

No 15, Jalan Mega 1/5,
Taman Perindustrian Nusa Cemerlang,
79200 Iskandar Puteri,
Johor, Malaysia

■ Revised Date

14 March 2024



Manufactured by

SAMJIN PHARM.CO.,LTD.

52, Jeyakongdan1-Gil, Hyangnam-Eup,
Hwaseong-Si, Gyeonggi-Do, Korea