

ACTIMIN

DESCRIPTION: ACTIMIN TABLET is a 7mm, round, blue tablet, scored on one side and marking 'd' on the other side.

COMPOSITION: Each tablet contains Triprolidine Hydrochloride 2.5mg and Pseudoephedrine Hydrochloride 60mg.

PHARMACOLOGY: Antihistamine and decongestant.

Tripolidine: Absorption and Fate: In general, antihistamines are readily absorbed from the gastro-intestinal tract, metabolised in the liver, and excreted usually mainly as metabolites in the urine.

Pseudoephedrine: Absorption and Fate: Pseudoephedrine is readily and completely absorbed from gastro-intestinal tract. It is resistant to metabolism by monoamine oxidase and is largely excreted unchanged in the urine. It has a half-life of several hours; elimination is enhanced and half-life accordingly shorter in acid urine.

INDICATIONS: Indicated in all disorders of the upper respiratory tract including the sinuses, and Eustachian tubes which are benefited by taking a potent decongestant and antihistamine. These include allergic (seasonal) rhinitis, vasomotor (perennial) rhinitis, hay fever, nasal congestion, and the common cold.

Actimin may be of value in the conservative management of secretory otitis media ("glue ear" and in helping to minimize gagging in patients undergoing dental procedures.

Because Actimin contains pseudoephedrine urinary retention may occasionally occur in male subjects where prostatic enlargement is present.

Fixed drug eruption due to pseudoephedrine, taking the form of erythematous nummular patches, and lichenoid skin eruption due to triprolidine have been reported but both these reactions should be regarded as rare events. Rarely, sleep disturbances and hallucinations have been reported.

RECOMMENDED DOSAGE:

Adults and Children over 12 Years: One tablet, 3 – 4 times as needed.

Children 6 – 12 years: ½ tablet 3 – 4 times a day as needed.

ROUTE OF ADMINISTRATION: Oral

CONTRAINDICATIONS: Actimin is contraindicated in individuals, who have previously exhibited intolerance to it or to pseudoephedrine or triprolidine. Actimin is contraindicated in patients who are taking or have taken monoamine oxidase inhibitors within the preceding two weeks. The concomitant use of pseudoephedrine and this type of product could occasionally cause a rise in blood pressure.

WARNING & PRECAUTIONS: WARNING: Actimin may cause drowsiness and the patient should be discouraged from driving a car or operating machinery. A warning to users against the concomitant use of alcohol or other centrally acting sedatives and Actimin is prudent.

PRECAUTIONS: Actimin should be used with caution in patients taking sympathomimetic agents such as decongestants, appetite suppressants and amphetamine-like psychostimulants, anti-hypertensive agents or other antihistamines. The effects of a single dose of Actimin on the blood pressure of these patients should be observed before recommending repeated or unsupervised treatment.

An extremely rare example of pseudoephedrine intolerance in a normotensive individual has been reported, following the ingestion of 60mg pseudoephedrine, which resulted in hypertension and unconsciousness.

As with other sympathomimetic agents and decongestants, Actimin should be used with caution in patients with prostatic enlargement or bladder dysfunction.

In severe hepatic or renal dysfunction, Actimin should be given at less than the usual recommended dose and the patient's response used as a guide to the dosage requirement for further administration.

Should not be used in new born or premature infants.

Antihistamines should not be used to treat lower respiratory tract symptoms including asthma.

In patients hypersensitive to triprolidine hydrochloride and sympathomimetic amines.

INTERACTION WITH OTHER MEDICATIONS: Concomitant use of Actimin with sympathomimetic agents, such as decongestants, appetite suppressants and amphetamine-like psychostimulants or with monoamine oxidase inhibitors which interfere with the catabolism of sympathomimetic amines, may occasionally cause a rise in blood pressure largely because of interaction with pseudoephedrine (see Contraindications and Precautions / Warnings).

The antibacterial agent, furazolidone, is known to cause a dose-related inhibition of monoamine oxidase. Although there are no reports of hypertensive crises caused by the concurrent administration of Actimin and furazolidone they should not be taken together.

Because of its pseudoephedrine content, Actimin may partially reverse the hypotensive action of drugs which interfere with sympathetic activity include bretylium, bethanidine, guanethidine, debrisoquine, methyldopa and beta adrenergic blocking agents.

Though objective evidence is lacking Actimin might, because it contains triprolidine, potentiate the sedative action of other antihistamines, alcohol and other sedatives of the central nervous system.

USE IN PREGNANCY & LACTATION: Use in Pregnancy: There has been no evidence that Actimin cause teratogenic effects. However, it is inadvisable to give any drug during the first trimester of pregnancy, unless circumstances demand.

Use in Lactation: Pseudoephedrine and triprolidine are excreted in breast milk in small amounts but the effect of this on breast fed infants is not known. It has been estimated that 0.5 to 0.7% of a single dose of pseudoephedrine ingested by a mother will be excreted in the breast milk over 24 hours.

SIDE EFFECTS: Actimin has the side effects of antihistamines. The most common effects are sedation, including inability to concentrate, lassitude, dizziness, hypotension, muscular weakness and incoordination.

Other side effects include gastro intestinal disturbances such as nausea, vomiting, diarrhoea or constipation, and epigastric pain. Antihistamines may also produce headache, blurred vision, tinnitus, elation or depression, irritability, nightmares, anorexia, difficulty in micturition, dryness of the mouth, tightness of the chest and tingling, heaviness and weakness of the hands. In infants and children, some antihistamines act as cerebral stimulants and symptoms of overdosage may include convulsions and hyperpyrexia.

Large doses of antihistamines may precipitate fits in epileptics.

SYMPTOMS AND TREATMENT OF OVERDOSE: Symptoms: The effects of acute toxicity from Actimin may include drowsiness, weakness, incoordination, difficulty with micturition, respiratory depression, hypotension, agitation, irritability, convulsions, hypertension, palpitations and tachycardia.

Treatment: Necessary measures should be taken to maintain and support respiration and circulation. Gastric lavage should be performed if indicated and convulsions controlled with diazepam. Catheterisation of the bladder may be necessary. Hypertension may be controlled with alpha adrenoceptor blocking drugs and tachycardia with beta adrenoceptor blocking drugs.

STORAGE CONDITIONS: Store below 30°C. Protect from light. Keep out of reach of children. *Jauhkan daripada kanak-kanak.*

PACK SIZE: Blister pack of 1000 tablets.

SHELF LIFE: Please refer to outer package. To consume within 2 months after opening the pouch.

PRODUCT REGISTRATION HOLDER AND MANUFACTURER:

DUOPHARMA (M) SDN BHD

Lot 2599 Jalan Seruling 59 Kawasan 3,
Taman Klang Jaya, 41200 Klang, Selangor, Malaysia.

1500011544 N.2