

Important Information. Please read carefully.

Asmaliv

Syrup

Sugar-free

Composition

Each 5 ml contains Salbutamol Sulphate equivalent to Salbutamol 2 mg. Sodium Benzoate 10mg/5ml as preservative.

Pharmacology

Salbutamol is readily absorbed from the gastro-intestinal tract. It is subjected to first-pass metabolism in the liver. After oral administration about 50% of the dose is excreted in the urine as an inactive sulphate conjugate, the rest being unchanged salbutamol. Its plasma half-life has been estimated to range from about 2 - 7 hours.

Indications

Salbutamol acts selectively on the bronchial muscle to relieve bronchospasm in patients with asthma and chronic bronchitis.

Dosage

In elderly patients or in those known to be unusually sensitive to beta-adrenergic stimulants drugs, it is advisable to initiate treatment with 2mg three or four times per day.

To be taken three or four times daily.

Children 2 to 6 years : 2.5 to 5 ml.

Children 6 to 12 years : 5 ml.

Over 12 years : 5 to 10 ml.

Precautions

Salbutamol should be used with caution in patients with hyperthyroidism. Great care is also needed in patients with cardiovascular disease such as ischaemic heart disease, arrhythmia or tachycardia, occlusive vascular disorders (including arteriosclerosis, hypertension, or aneurysms) and angina pectoris. Care is also required when Salbutamol is given to patients with diabetes mellitus, elderly patients and closed-angle glaucoma.

Usage During Pregnancy

Administration of the drug during pregnancy should only be considered if the expected benefit to the mother is greater than any possible risk to the foetus or neonate.

Interaction

Salbutamol should be avoided or used with caution in patients undergoing anaesthesia with cyclopropane, halothane, or other halogenated anaesthetics. An increased risk of arrhythmia may also occur if Salbutamol

is given to patients receiving cardiac glycosides, quinidine, or tricyclic antidepressants.

Propranolol and other beta-adrenoceptor blocking agents antagonise the effects of Salbutamol. Hypokalaemic effects caused by Salbutamol may also be enhanced by concomitant administration of aminophylline or other xanthines, and diuretic therapy.

Contraindications

It is contraindicated in patients with a history of hypersensitivity to any of their components. It is not indicated for the prevention of premature labour associated with toxemia of pregnancy or antepartum haemorrhage, nor should it be used for threatened abortion during the first & second trimesters of pregnancy.

Side Effects

Mild tremor (usually with the hands) may occur in some patients. Other side effects that may occur include headache, nervous tension, peripheral vasodilatation and tachycardia.

Hypersensitivity reactions including angioedema, urticaria, bronchospasm, hypotension and serious hypokalaemia.

Overdosage & Treatment

Symptoms: Symptoms of overdosage may include the adverse effects described above. Significant tachycardia, muscle tremor and superficial vasodilatation are common.

Treatment: The stomach should be emptied by gastric lavage. The specific antidote is a selective beta-1 adrenergic receptor blocking agent given by intravenous injection. However, beta-blockers should be used cautiously in patients with a history of bronchospasm.

Shelf life

2.5 years from date of manufacture.

Presentation

Clear, colourless, kiwi flavoured syrup in amber PET bottle of 50ml, 60ml, 90ml, 100ml and 120ml.

Storage

Store below 25°C.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

MAL 19890165A

SIN 5105P

HK 36075



Xepa^{sp}

Manufactured by:

Xepa-Soul Pattinson (Malaysia) Sdn. Bhd.

1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia.