



## PRIME'S CHLORPHENIRAMINE SYRUP 4MG / 5ML

Description: Red coloured and strawberry flavoured syrup.

### Contents:

|                     |                          |          |
|---------------------|--------------------------|----------|
| Each 5ml contains : | Chlorpheniramine Maleate | 4mg      |
|                     | Methylparaben            | 0.1%w/v  |
|                     | Propylparaben            | 0.01%w/v |

### Indications:

For treatment of perennial and seasonal allergic rhinitis, vasomotor rhinitis, allergic conjunctivitis, allergic skin manifestation of urticaria and angioedema.

### Pharmacology:

Chlorpheniramine acts by competing with histamine for H<sub>1</sub>-receptor sites on effector cells, thereby preventing responses mediated by histamine. Its muscarinic actions provide a drying effect on the nasal mucosa. It is readily absorbed orally, metabolised in the liver and excreted in the urine.

### Adverse Effects:

Drowsiness, dryness of mouth nose and throat, gastrointestinal disturbances, and skin rash may occur.

### Precautions

- Caution in patients with glaucoma, acute asthma and renal function impairment.
- Caution when driving or doing jobs requiring alertness because of possible drowsiness. Avoid alcoholic beverages.
- Safety for use in pregnancy and lactation has not been established.

### Warning:

- When used for treatment of cough and cold;
- not to be used in children less than 2 years of age.
- To be used with caution and on doctor's or pharmacist's advice in children 2 to 6 years of age.

### Contraindications:

- Avoid in patients known to be hypersensitive to antihistamines.
- Use is not recommended in nursing mothers.

### Drug Interactions:

- Concurrent use of alcohol, barbiturates, tricyclic antidepressants, narcotic analgesics, or atropine may potentiate the effects of either these medications or antihistamines.
- Co-administration of monoamine oxidase (MAO) inhibitors may prolong and intensify the antimuscarinic effects of antihistamines.

### Overdosage:

Clinical signs include severe drowsiness, unsteadiness, hallucinations, seizures, and respiratory depression. Since there is no specific antidote for overdose with antihistamines, treatment is symptomatic and supportive.

### Dosage and Administration:

|          |   |                                                |
|----------|---|------------------------------------------------|
| Adults   | : | - Oral, 1 teaspoonful three times a day.       |
| Children | : | - Oral, 1/2 - 1 teaspoonful three times a day. |
|          | : | - Oral, 1/2 teaspoonful three times a day.     |

Storage : Keep container tightly closed. Store upright in cool, dry place below 30°C.  
Protect from light. Keep out of reach of children.

Presentation/ Packing : Syrup 4mg / 5ml x 60ml

Manufactured & Distributed by: **PRIME PHARMACEUTICAL SDN. BHD.**  
1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,  
14000 Bukit Mertajam, Pulau Pinang, Malaysia.