

Important information. Please read carefully.



Fenfedrin™

Capsules

Relieves nasal congestion in common colds and allergic rhinitis (eg. hayfever).

Composition

Each capsule contains

Chlorpheniramine Maleate	4mg
Pseudoephedrine Hydrochloride	60mg

Pharmacodynamics

Chlorpheniramine Maleate belongs to the alkylamine antihistamines, which are among the most active H₁-blockers and are generally effective in relatively low doses. It is used to reduce rhinorrhoea and sneezing.

Pseudoephedrine is a sympathomimetic agent with direct and indirect effects on adrenergic receptors. It is said to have less pressor activity and CNS effects than ephedrine of which it is a stereoisomer. Pseudoephedrine hydrochloride is used as a nasal and bronchial decongestant.

Both active ingredients are readily absorbed from the gastro-intestinal tract.

Pseudoephedrine is resistant to metabolism by monoamine oxidase and up to about 90% of a dose in 24 hours is largely excreted unchanged in the urine. Less than 1% is excreted as nor-pseudoephedrine which is pH dependent. The half-life of several hours is reduced in acid urine. Chlorpheniramine is extensively metabolised to polar and non-polar metabolites. The excretion of the unchanged drug is increased by acidification of the urine and increased flow of urine. After daily oral administration the forms excreted are about 20% unchanged drug, 20% as monodesmethylchlorpheniramine, 5% as didesmethylchlorpheniramine. Less than 1% is excreted through the faeces.

Pharmacokinetics

Both active ingredients are readily absorbed from the gastro-intestinal tract.

Pseudoephedrine is resistant to metabolism by monoamine oxidase and up to about 90% of a dose in 24 hours is largely excreted unchanged in the urine. Less than 1% is excreted as nor-pseudoephedrine which is pH dependent. The half-life of several hours is reduced in acid urine.

Chlorpheniramine is extensively metabolised to polar and non-polar metabolites. The excretion of the unchanged drug is increased by acidification of the urine and increased flow of urine. After daily oral administration the forms excreted are about 20% unchanged drug, 20% as monodesmethylchlorpheniramine, 5% as didesmethylchlorpheniramine. Less than 1% is excreted through the faeces.

Indication

Fenfedrin alleviates nasal congestions in common colds and helps clear nasal passages. It also provides temporary relief of runny nose, sneezing, itching of the nose which may occur in allergic rhinitis such as hayfever.

Dosage

To be administered orally. Unless otherwise prescribed by the physician:

Adults: One capsule every four to six hours.

Not recommended for children under 12 years of age.

Contraindications

This medicine is contraindicated in patients taking monoamine oxidase inhibitors and within two weeks from stopping treatment. It is also contraindicated in patients with known hypersensitivity to chlorpheniramine and pseudoephedrine.

Precautions

This may cause drowsiness. If affected do not drive motor vehicles or operate machinery. Do not drink alcohol while taking this medicine.

Use with caution in patients with asthma, glaucoma, epilepsy, prostatic hypertrophy, peptic ulcer, pyloroduodenal or bladder neck obstruction, hyperthyroidism, heart disease or hypertension. It is not advisable for use in pregnancy except under the advice and supervision of a physician.

Drug Interaction

It should not be used for patients taking or have taken antidepressant therapy containing monoamine oxidase inhibitors within the two preceding weeks.

Pregnancy and Lactation

It is not advisable for use in pregnancy except under the advice and supervision of a physician.

Side Effects

Side effects that may occur include drowsiness, headache, dry mouth, allergic reactions such as rashes, gastrointestinal disturbances, tightness of chest and in rare cases, certain blood disorders.

Overdosage and Treatment

Paranoid psychosis, delusions, hallucinations may occur with overdosage due to the pseudoephedrine component. With chlorpheniramine, convulsions, hyperpyrexia, stimulation effects such as tachycardia, nervousness and convulsions may occur. Treat nervous stimulation and convulsions with a sedative such as intramuscular diazepam. In severe cases, stomach should be emptied by aspiration and gastric lavage. Elimination of pseudoephedrine may be assisted by acidification of the urine with ammonium chloride.

Shelf life

The expiry date is indicated on the packaging.

Presentation

Size 2, Grey/White hard gelatin capsules with 'XSP' printed in dispensing packs of 7's and 35's.

Storage

Store in a cool, dry place below 30°C.

KEEP OUT OF REACH OF CHILDREN/JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date : 17-Nov-2021

Manufactured and Product Registration Holder (in Malaysia)

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia