



## PRIME'S PREDNISOLONE SYRUP 2.5mg/5ml

Red coloured, cherry flavoured syrup.

### Contents:

Prednisolone 2.5mg/ 5ml  
Methylparaben 0.1% w/v  
Propylparaben 0.01% w/v

### Indications:

Corticosteroid indicated conditions include allergic disorders such as bronchial asthma and allergic skin reactions; blood disorders, collagen and rheumatic disorders; connective tissue disorders; gastrointestinal disorders such as inflammatory bowel disease.

### Pharmacology:

Prednisolone, a synthetic glucocorticoid, is used for its anti-inflammatory action in the treatment of conditions in which routine systemic corticosteroid therapy is indicated, however, it has a weaker sodium-retaining effect than cortisone. It is readily absorbed orally and has a plasma half-life of 2 to 3 hours. It undergoes hepatic metabolism and renal excretion.

### Adverse effects:

Hypokalemia, osteoporosis, Cushing's syndrome, acne or other skin problems, gastrointestinal bleeding, endocrine imbalance, hyperglycaemia, and increased susceptibility to infections.

Sodium retention electrolyte imbalance, and edema are less pronounced.

### Precautions:

- Caution in patients with congestive heart failure, diabetes mellitus, renal or hepatic function impairment hypertension, diverticulitis, myasthenia gravis, hypothyroidism, infectious diseases, tuberculosis and in elderly persons.
- Children on prolonged therapy should be closely observed.
- Treatment should not be discontinued without the advice of the prescriber.
- Safety for use in pregnancy and lactation has not been established.
- **Scleroderma renal crisis**-Caution is required in patients with systemic sclerosis because of an increased incidence of (possibly fatal) scleroderma renal crisis with hypertension and decreased urinary output observed with a daily dose of 15mg or more prednisolone.

### Contraindications:

Not recommended for patient with peptic ulcer, osteoporosis, psychoses or severe psychoneuroses, acute infections including herpes zoster and herpes simplex ulceration of the eye, and vaccination with live vaccine.

### Drug interactions:

- Co-administration of barbiturates, phenytoin, or rifampicin, may reduce the effects of corticosteroids.
- Response to anticoagulants and salicylates may be reduced.
- Concurrent use of potassium-depleting diuretics may cause excessive potassium loss.

### Dosage and administration:

Adults : Oral, 5 to 60mg a day, the dosage being adjusted according to patient response.  
Children : As prescribed by the physician.

### Overdosage

Adverse effects are almost always signs of overdosage. Dosage should be reduced or the drug slowly withdrawn. Symptomatic and supportive therapy is recommended.

Storage conditions : Store in a cool dry place below 30°C. Protect from light. Keep out of reach of children.

Presentation/ Packing: Syrup 2.5mg/ 5ml X 60ml

Shelf life : 2 years

Manufactured & : **PRIME PHARMACEUTICAL SDN. BHD.**

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,  
14000 Bukit Mertajam, Penang, Malaysia.