



TRIPROLIDINE-P SYRUP

Description: Yellow coloured, lemon flavoured syrup.

Contents:

Each 5ml contains :	Tripolidine HCl B.P.	1.25mg
	Pseudoephedrine HCl B.P.	30mg
Preservatives :	Methylparaben	0.1%
	Propylparaben	0.01%

Indications:

Nasal and respiratory congestion, common cold, acute sinusitis, allergic rhinitis.

Pharmacology:

Tripolidine HCl, a potent antihistamine, provides a driving effect on the nasal mucosa by means of its antimuscarinic actions. Pseudoephedrine, a sympathomimetic amine, acts on alpha-adrenergic receptors in the mucosa of the respiratory tract to produce vasoconstriction which reduces the swelling associated with inflammation of the mucous membranes lining the nasal passage. Oral absorption is rapid with onset of action about 15 to 30 minutes.

Adverse Effects:

Drowsiness, lassitude, dizziness, nausea, vomiting, epigastric pain, blurred vision, confusion, difficulty in micturition, palpitations, headache, tinnitus, dryness of mouth, anorexia, and restlessness may occur occasionally.

Precautions:

- Caution in patients with cardiovascular disease, glaucoma, hypertension, hyperthyroidism and prostatic hypertrophy.
- Caution when driving or doing jobs requiring alertness because of possible drowsiness. Avoid alcoholic beverages.
- Safety for use in pregnancy has not been established.

Warning:

- Not to be used in children less than 2 years of age.
- To be used with caution and on doctor's or pharmacist's advice in children 2 to 6 years of age.

Contraindications:

- Avoid in patients intolerant to tripolidine or pseudoephedrine.
- Use is not recommended in nursing mothers.
- It should not be given to patients who are being treated with a monoamine-oxidase (MAO) inhibitor or within 2 weeks of discontinuation such treatment.

Drug Interactions:

- Co-administration of atropine or other antimuscarinics may potentiate the muscarinic effects.
- Concurrent use of tricyclic antidepressants may potentiate the CNS-depressant effects of either medications.

Dosage and Administration:

Adults	-	Oral, 2 teaspoonfuls three or four times a day as needed
Children	- 6 to 12 years	- Oral, 1 teaspoonful three or four times a day.
	- 2 to 5 years	- Oral, 1/2 teaspoonful three times a day.

Overdosage:

Clinical signs include unsteadiness, flushing, shortness of breath, hallucinations, drowsiness, seizures, severe dryness of mouth, nose or throat, continuing headache and unusually slow or fast heartbeat.

Treatment consists of emesis or gastric lavage, if appropriate, and the necessary symptomatic and supportive therapy.

Storage : Store in a cool, dry place, below 30°C. Protect from light.
Keep out of reach of children.

Presentation/ Packing : Syrup x 90ml

Manufactured & Distributed by : **PRIME PHARMACEUTICAL SDN. BHD.**
1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

Further information can be obtained from your doctor or pharmacist

13-04-2017