

DECAN

INJECTION

**Ingredients:** Each ml. contains:

Dexamethasone Sodium Phosphate	2.2mg
(Equivalent to Dexamethasone Phosphate 2.0mg)	
Preservative:	
Methylparaben.....	1.5mg
Propylparaben.....	0.2mg

Pharmacodynamics:

Dexamethasone Phosphate can decrease or prevent tissue responses to inflammatory processes, thereby reducing development of symptoms of inflammation without affecting the underlying cause. Dexamethasone inhibit accumulation of inflammatory cells, including macrophages and leukocytes at sites of inflammation. They also inhibit phagocytosis, lysosomal enzyme release and synthesis and/or release several chemical mediators of inflammation. Although the exact mechanisms are not completely understood, actions which may contribute significantly to these effects include blockade of the action of macrophage inhibitory factor (MIF), leading to inhibition of macrophage localization, reduction of dilatation and permeability of inflamed capillaries and reduction of leukocyte adherence to the capillary endothelium, leading to inhibition of both leukocyte migration and edema formation ; and increased synthesis of lipomodulin (Macroscortin), an inhibitor of phospholipase A₂-mediated arachidonic acid release for membrane phospholipids with subsequent inhibition of the synthesis of arachidonic acid-derived mediators of inflammation (Prostaglandins, thromboxanes, and leukotrienes). Immunosuppressant actions may also contribute significantly to the anti-inflammatory effect.

Pharmacokinetics:

1. Dexamethasone Phosphate is a synthetic glucocorticoid and has a potent anti-inflammatory property. It is indicated for the treatment of endocrine disorders, rheumatic disorder, collagen diseases, dermatologic diseases, and allergic states etc.

2. Dexamethasone Phosphate is rapidly absorbed after intramuscular route. It is metabolized in the liver and excreted in the urine, mainly as inactive metabolites.

Route of Administration: I.V or I.M**Indications:**

Treatment of acute or chronic rheumatism, rheumatoid arthritis, acute gout arthritis, myalgia, eczematosis, allergic dermatitis, and acute or chronic allergic and inflammatory skin disorders.

Dosage and Administration:

- For the treatment of acute adrenocortical insufficiency, bronchial asthma, serious infections, acute drug poisoning.
For adults: it is given by intravenous or intramuscular route. Injection in a dose of 2-10mg, several times a day. Do not exceed 80mg per day.
- For the treatment of osteoarthritis, rheumatoid arthritis, acute gouty arthritis:
Adult: a. for genu articular by intra-articular injection, 4-6mg each time.
b. for manus articular, 0.8-1 mg each time.
- For the treatment of myositis fibrositis sciatic neuralgia:
For adult: 2-6 mg by soft tissue injection.
- For the treatment of contact dermatitis, infant eczema , dermatopolynuritis and psoriasis:
For adult: 0.5mg by subcutaneous injection, diluted 5 times with normal saline solution before administration.
To be used only by a physician.

Symptoms And Treatment of Overdose:

There is no specific antidote for its overdose, therefore, management of the patient consist of symptomatic and supportive therapy.

Side effects:

- The prolonged therapy may result in suppression of pituitary-adrenal function and the principal complications resulting from prolonged therapy with Dexamethasone Phosphate are fluid and electrolyte disturbance; hyperglycemia and glycosuria; increased susceptibility to infections, including tuberculosis; peptic ulcer, which may bleed or perforate; osteoporosis; a characteristic myopathy; behavioral disturbances; posterior subcapsular cataracts; arrest of growth ; and Cushing's habitus, consisting of "moon face", "buffalo hump", enlargement of supraclavicular fat pads, "central obesity", striae, ecchymoses, acne, and hirsutism.
- Acute adrenal insufficiency results from too rapid withdrawal of corticosteroids after prolonged therapy. Dexamethasone withdrawal syndrome, consisting of fever, myalgia, arthralgia and malaise.

Contraindications:

Dexamethasone Phosphate is contraindicated in patients with systemic fungal infections and those who have shown hypersensitivity to any component of this product.

Warning and Precautions:

- The adverse effects of dexamethasone are nearly always due to their use in excess of normal physiological requirement. They should be treated symptomatically, where possible the dosage being reduced or the drugs slowly withdrawn.
- Dexamethasone may mask some signs of infection and new infections may appear during its use. There may be decreased resistance and inability to localize infection when dexamethasone is used.
- Dexamethasone should be used cautiously in patients with ocular herpes simplex, psychic derangements and gastric, duodenal ulcer.

Interaction with Other Medicaments:

None has been mentioned

Pregnancy and Lactation:

The use of dexamethasone in pregnancy, nursing mothers, or women of childbearing potential requires that possible benefits of the drug be weighed against the potential hazards, to the mother and embryo or fetus.

Storage Condition:

Store away from heat and direct light. Store at temperature below 30°C.

Shelf-life:

3 years

Product Description:

A clear, colorless solution.

Packaging:

Box of 2ml x 10 ampoule and 2mlx 100 ampoule.



Manufacturer:

YUNG SHIN PHARMACEUTICAL IND. CO., LTD.

Taichung Youth Factory
No. 27, Kung Chiu Rd., Tachia, Taichung, Taiwan, R.O.C.

Importer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD.

Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.

Ordering Line: 1 800 88 3027 Product Info: 1 800 88 3679



2146134 Date of Revision: 01 Dec 16