



PRISOLONE TABLET 5mg

Description: White, square, bevel-edged top scored tablet.

PREDNISOLONE SYRUP 2.5mg/5ml

Description: Red coloured, cherry flavoured syrup.

Contents:

Prisolone Tablet	- Prednisolone 5mg/ tablet
Prednisolone Syrup	- Prednisolone 2.5mg/ 5ml
	Methylparaben 0.1% w/v
	Propylparaben 0.01% w/v

Indications:

Corticosteroid indicated conditions include allergic disorders such as bronchial asthma and allergic skin reactions; blood disorders, collagen and rheumatic disorders; connective tissue disorders; gastrointestinal disorders such as inflammatory bowel disease.

Pharmacology:

Prednisolone, a synthetic glucocorticoid, is used for its anti-inflammatory action in the treatment of conditions in which routine systemic corticosteroid therapy is indicated, however, it has a weaker sodium-retaining effect than cortisone. It is readily absorbed orally and has a plasma half-life of 2 to 3 hours. It undergoes hepatic metabolism and renal excretion.

Adverse effects:

Hypokalemia, osteoporosis, Cushing's syndrome, acne or other skin problems, gastrointestinal bleeding, endocrine imbalance, hyperglycaemia, and increased susceptibility to infections.

Sodium retention electrolyte imbalance, and edema are less pronounced.

Precautions:

- Caution in patients with congestive heart failure, diabetes mellitus, renal or hepatic function impairment hypertension, diverticulitis, myasthenia gravis, hypothyroidism, infectious diseases, tuberculosis and in elderly persons.
- Children on prolonged therapy should be closely observed.
- Treatment should not be discontinued without the advice of the prescriber.
- Safety for use in pregnancy and lactation has not been established.

Scleroderma renal crisis:

- Caution is required in patients with systemic sclerosis because of an increased incidence of (possibly fatal) scleroderma renal crisis with hypertension and decreased urinary output observed with a daily dose of 15 mg or more prednisolone.

Contraindications:

Not recommended for patient with peptic ulcer, osteoporosis, psychoses or severe psychoneuroses, acute infections including herpes zoster and herpes simplex ulceration of the eye, and vaccination with live vaccine.

Drug interactions:

- Co-administration of barbiturates, phenytoin, or rifampicin, may reduce the effects of corticosteroids.
- Response to anticoagulants and salicylates may be reduced.
- Concurrent use of potassium-depleting diuretics may cause excessive potassium loss.

Dosage and Administration:

Adults	: Oral, 5 to 60mg a day, the dosage being adjusted according to patient response.
Children	: As prescribed by the physician.

Overdosage:

Adverse effects are almost always signs of overdosage. Dosage should be reduced or the drug slowly withdrawn. Symptomatic and supportive therapy is recommended.

Storage conditions : Store in cool dry place below 30°C. Protect from light.
Keep away from children.

Presentation/ Packing: Tablet 5mg x 100x10's; Syrup 2.5mg/ 5ml X 60ml

Shelf life : Tablet : 3 years Syrup:2 years

Product registration : **PRIME PHARMACEUTICAL SDN. BHD.**

holder & manufacturer 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.