



Xepa^{SP}

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia.

Important Information. Please read carefully.

ZarinTM

Cream

COMPOSITION

Contains Miconazole Nitrate 2%w/w in a non-greasy, non-staining cream basis. Benzoic Acid 0.15% w/w as preservative.

PHARMACODYNAMICS

Miconazole is an imidazole antifungal agent that interferes with ergosterol synthesis and therefore alters the permeability of the cell membrane of sensitive fungi. Miconazole is active against the common dermatophytes *Trichophyton rubrum*, *T. mentagrophytes* and *Epidermophyton*; the yeast *Candida albicans*; and the organism responsible for tinea versicolor, *Malassezia furfur*.

PHARMACOKINETICS

There is little absorption through skin or mucous membranes when miconazole nitrate is applied topically.

Miconazole is metabolized in the liver to inactive metabolites.

The half life of miconazole is about 24 hours.

INDICATIONS

Zarin is indicated for the topical treatment of skin infections due to susceptible fungi (dermatophytes and yeasts) such as tinea pedis (athlete's foot), tinea cruris, tinea corporis (ringworm), tinea versicolor (white spots), and candidiasis.

DOSAGE AND ADMINISTRATION

Apply a thin layer of Zarin sufficient to cover the affected areas and rub well into the skin until the cream has been completely absorbed. The cream should be applied once or twice a day, in the morning and evening, and continued without interruption until the symptoms have been completely disappeared (usually 2 to 5 weeks, depending on the infecting organism and the site of infection).

CONTRAINDICATIONS

Zarin is contraindicated in patients known to be hypersensitive to imidazole antifungal agents or any components of the medication.

WARNINGS AND PRECAUTIONS

Discontinue treatment if sensitisation or irritation occurs during use.

DO NOT USE IN THE EYES.

It is advisable to consult a doctor if no improvement is observed in 4 weeks in the case of 'athlete's foot' or 2 weeks in 'jock itch'.

In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see *Interactions*).

INTERACTIONS

Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and the anticoagulant effect should be monitored (see *Warnings and Precautions*).

SIDE EFFECTS

Zarin is usually well tolerated. However, local irritation and sensitivity reactions may occasionally occur.

OVERDOSAGE AND TREATMENT

Accidental ingestion by mouth may cause mild gastrointestinal disturbances. Symptomatic and supportive treatment may be of value.

SHELF-LIFE

The expiry date is indicated on the packaging.

PRESENTATION

White, smooth, homogeneous, non-greasy and water-miscible cream in tubes of 15g.

INGREDIENTS

Miconazole Nitrate, Benzoic Acid, Cetyl Alcohol, Liquid Paraffin, Propylene Glycol, Sorbitan Monostearate, Stearic Acid, Polysorbate 80, Purified Water.

STORAGE

Store below 30°C.

KEEP OUT OF REACH OF CHILDREN**JAUHI DARI KANAK-KANAK**

For further information, please consult your pharmacist or physician.

Revision Date : 01-Dec-2017

Manufacturer and Product Registration Holder:

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia.