

BECARIN CREAM



DESCRIPTION

BECARIN CREAM contains 2%w/w Miconazole Nitrate in a white cream base.

Preservatives:

Methylparaben 0.32%w/w, Propylparaben 0.048%w/w.

PHARMACODYNAMICS

Miconazole is fungistatic, inhibits biosynthesis of ergosterol or other sterols, damaging the fungal cell wall membrane and altering permeability; as a result, loss of essential intracellular elements may occur, also inhibits biosynthesis of triglycerides and phospholipids by fungi; in addition, inhibits oxidative and peroxidative enzyme activity; resulting in intracellular buildup of toxic concentrations of hydrogen peroxides which may contribute to deterioration of subcellular organelles and cellular necrosis. In *Candida albicans*, inhibits transformation of blastospores into invasive mycelial form.

PHARMACOKINETICS

Miconazole is incompletely absorbed from the gastro-intestinal tract. Penetration into cerebrospinal fluid and sputum is poor but miconazole infuses well into infected joints. Over 90% reported to be bound to plasma proteins. Miconazole is metabolised in the liver to inactive metabolites. There is little absorption through skin and mucous membranes when miconazole nitrate is applied topically.

INDICATIONS

Treatment of fungal infections of skin and nails including cutaneous candidiasis, tinea corporis, tinea cruris, tinea pedis and tinea versicolor (pityriasis versicolor).

APPLICATION

Apply to the affected areas two or three times daily continuing for fourteen days after lesions have healed.

CONTRAINDICATIONS

Sensitivity to topical miconazole.

Route of administration: Topical.

PRECAUTIONS/WARNING

Do not use on children under 2 years of age except under the advice and supervision of a doctor. If irritation occurs or if there is no improvement within 4 weeks, discontinue use and consult a doctor or pharmacist.

In patients with a warfarin, caution should be exercised and the anticoagulant effect should be monitored. (See Interactions)

INTERACTIONS

Miconazole administered systematically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and the anticoagulant effect should be monitored (see Warnings and Precautions)

SIDE EFFECTS/ADVERSE REACTIONS

Occasional skin irritation or sensitivity.

STORAGE CONDITIONS AND USER INSTRUCTIONS

Keep container tightly closed.

Store in a dry place below 30°C.

Protect from light.

Keep out of reach of children.

Jauhi daripada kanak-kanak.

Shelf life : Please refer to outer package.

PRESENTATION

BECARIN CREAM is supplied in 15g and 30g tubes.

Product Registration Holder:

Duopharma Manufacturing (Bangi) Sdn.Bhd.

Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13,
Bangi Industrial Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.

Manufacturer:

Duopharma Manufacturing (Bangi) Sdn.Bhd.

Lot No. 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate,
43650 Bandar Baru Bangi, Selangor, Malaysia.



1500008648 N.1
Revision Date: May 2019

BECARIN CREAM

16.9mm(w) x 16.3mm (h)



Black 100%