



MICONAZOLE CREAM

MAL19900255AZ

COMPOSITION:

Miconazole Nitrate (micronised) 2% w/w

Chlorocresol 0.1% w/w as preservative.

PRESENTATION:

A white, water-miscible cream.

INDICATIONS:

Miconazole is used topically in the management of superficial candidiasis and fungal skin infections, such as; mucocutaneous candidiasis, dermatophytosis, and pityriasis versicolor.

PHARMACOLOGY:*Mechanism of Action:*

Miconazole is an imidazole antifungal agent with some antibacterial activity. It has some activity against *Aspergillus spp.*, *Cryptococcus neoformans*, *Pseudallescheria boydii*, and some Gram-positive bacteria including *staphylococci* and *streptococci*.

Pharmacokinetics:

Miconazole is absorbed from the gastrointestinal tract. Up to 20% of a dose given by mouth is excreted mainly as metabolites in the urine over 4 days. Up to 55% is excreted unchanged in the faeces over 6 days. There is little absorption through skin or mucous membranes.

DIRECTION FOR USE:

For external use only.

Apply cream (according to the size of lesion) to the lesion twice daily. Rub gently until the cream has been completely absorbed by the skin. The treatment should be continued without interruption until the lesion has completely disappeared (usually after 2 to 5 weeks). Continue treatment for at least 1 week after the lesion has completely disappeared to prevent relapse. If irritation occurs or if there is no improvement in 4 weeks (for athlete's foot) or 2 weeks (for jock itch), discontinue and consult doctor or pharmacist.

CONTRAINDICATION:

No known contraindication.

PRECAUTION:

Do not use in children under 2 years of age except under the advice and supervision of a doctor. Avoid contact with eyes. In patients on

warfarin, caution should be exercised and the anticoagulant effect should be monitored.

DRUG INTERACTIONS:

Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systematic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and the anticoagulant effect should be monitored.

SIDE-EFFECTS:

Local irritation and sensitivity reactions may occur when used topically.

OVERDOSAGE & TREATMENT:

Topical application of Miconazole Nitrate is well tolerated as there is little absorption through the skin. No known systemic toxicity of the drug has been reported when applied topically. In cases of accidental ingestion of large quantities of the product, an appropriate method of gastric emptying may be used if considered necessary.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK****PACK QUANTITIES:**

Available in aluminium tube of 5gm and 15gm.

Further information can be obtained from pharmacist, physician or the manufacturer.

Product Registration Holder & Manufactured By :



Kotra Pharma (M) Sdn. Bhd.

No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision : 01/03/24