

PRODUCT LITERATURE

FUNGICIDE CREAM 2%W/W

Each 100 g contains

Miconazole Nitrate 2 g

Preservative

Chlorocresol 0.1% w/w

Description

White coloured odourless soft cream.

Pharmacodynamics properties

Miconazole nitrate is an imidazole antifungal agent and may act by interfering with the permeability of the fungal cell membrane. It possesses a wide antifungal spectrum and has some antibacterial activity.

Pharmacokinetics properties

Absorption: There is little absorption through skin or mucous membranes when miconazole nitrate is applied topically.

Distribution: Absorbed miconazole is bound to plasma proteins (88.2%) and red blood cells (10.6%).

Metabolism and Excretion: The small amount of miconazole that is absorbed is eliminated predominantly in faeces as both unchanged drug and metabolites.

Indication

Treatment of fungal infections of the skin and nails including cutaneous candidiasis, tinea corporis, tinea cruris, tinea pedis and tinea versicolor (pityriasis versicolor)

Recommended Dosage

Apply a thin layer on the affected part 2 or 3 times a day until complete resolution of lesion

Route of Administration

Topical

Contraindications

Hypersensitivity to the active substances, other imidazole derivatives or to any of the excipients.

Warnings and Precautions

Fungicide Cream must not come into contact with the mucosa of the eyes.

Severe hypersensitivity reactions, including anaphylaxis and angioedema, have been reported during treatment with other miconazole topical formulations. If a reaction suggesting hypersensitivity or irritation should occur, the treatment should be discontinued.

In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see Interactions).

Interactions with Other Medicaments

Miconazole administered systemically is known to inhibit CYP3A4/2C9. Due to the limited systemic availability after topical application, clinically relevant interactions are rare. However, in patients on oral anticoagulants, such as warfarin, caution should be exercised and anticoagulant effect should be monitored. (see Warnings and Special Precautions).

Pregnancy and Lactation:

Pregnancy

In animals miconazole nitrate has shown no teratogenic effects but is foetotoxic at high oral doses. Only small amounts of miconazole nitrate are absorbed following topical administration. However, as with other imidazoles, miconazole nitrate should be used with caution during pregnancy.

Breast-feeding

Topically applied miconazole is minimally absorbed into the systemic circulation, and it is not known whether miconazole is excreted in human breast milk. Caution should be exercised when using topically applied miconazole products during lactation.

Side Effects / Adverse Reactions

This medication may cause irritation in some susceptible patients. If blistering, burning, redness, skin rash or other signs of skin irritation is experienced, then discontinue this medication and consult a doctor / pharmacist.

Overdose and Treatment

Symptoms

Cutaneous use: Excessive use can result in skin irritation, which usually disappears after discontinuation of therapy.

Treatment

Fungicide cream is intended for cutaneous use, not for oral use. If accidental ingestion of large quantities of the product occurs, use appropriate supportive care

Storage

Store below 30°C in a dry place, protected from direct light.

Keep away from reach of children.

Shelf Life

2 years from date of manufacture

Packing

Packed in plastic container of 5 g, 10 g and 50 g.

Name and Address of Product Registration Holder / Distributor

ZONTRON PHARMACEUTICALS SDN. BHD.
(445695-T)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia.

Name and Address of Product Manufacturer :

TERAPUTICS SDN. BHD. (590500-W)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia.

Malaysian Drug Registration No.: MAL19900274ACZ

Date of revision: 10/08/2021