

PACKAGE INSERT



MICONAZOLE CREAM 2%W/W

Composition:
Miconazole nitrate 2% w/w

Product Description: Smooth white cream.

Pharmacodynamics:

Miconazole nitrate has anti-fungal activity against *Trichophyton rubrum*, *Trichophyton mentagrophytes* and *Epidermophyton floccosum*, *Candida albicans* and organism responsible for tinea versicolor. It may act by interfering with the permeability of the fungal cell membrane. It also possesses some anti-bacterial activity against gram-positive bacteria including some staphylococci and streptococci.

Pharmacokinetics:

After topical application to the skin of normal subjects, systemic absorption of miconazole nitrate is extremely low. Although most of the applied drug remains on the skin surface, drug concentrations were found in the stratum corneum which, by far, exceeded the minimum inhibitory concentration for dermatophytes. Inhibitory concentrations were achieved in the epidermis and as deep as the middle region of the dermis.

Indication:

For topical application in the treatment of tinea pedis, tinea cruris and tinea corporis caused by *Trichophyton rubrum*, *Trichophyton mentagrophytes* and *Epidermophyton floccosum*, in the treatment of cutaneous candidiasis and in the treatment of tinea versicolor.

Recommended Dosage:

Sufficient cream should be applied to cover affected areas twice daily (morning and evening) in patients with tinea pedis, tinea cruris, tinea corporis and cutaneous candidiasis, and once daily in patients with tinea versicolor. Candidal infections and tinea cruris and corporis should be treated for two weeks, and tinea pedis for one month in order to reduce the possibility of recurrence.

Route of Administration: Topical

Contraindications:

It is contraindicated in patients with a history of hypersensitivity.

Warnings and Precautions:

1. Do not use in children under 2 years of age except under the advice and supervision of a doctor.
2. If irritation occurs or if there is no improvement within 4 weeks, discontinue use and consult a doctor or pharmacist.

In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see Interactions).

Interactions with Other Medicaments:

Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and the anticoagulant effect should be monitored (see Warnings and Precautions).

Pregnancy and Lactation:

Caution is recommended during pregnancy and lactation period. Treatment of large surfaces and the application under occlusive dressing should be avoided during that time.

Side Effects:

Reports of irritation, burning and maceration have been associated with the use of miconazole nitrate cream. Contact dermatitis has also been reported.

Symptoms and Treatment of Overdose:

Overdosage of miconazole nitrate arising from the systemic absorption from topical application has not been reported to date.

Effects on Ability to Drive and Use Machine: Not Applicable.

Storage Conditions: Store at or below 30°C.

Pack Size: A tube of 15g and 30g.

A box of 50x15g/ tube.

Pack Size (export only): A jar of 400g.

Shelf-life: 3 years.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer & Product Registration Holder:

SUNWARD PHARMACEUTICAL SDN. BHD.

No. 9, 11 & 17, Jalan Kempas 4,
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.

EL213A-R8
Revision Date: 07/10/2025