



MICOZOL CREAM 2%W/W

Description: White colour cream

Contents

Miconazole Nitrate 2% w/w
Chlorocresol 0.1% w/w

Indications

For treatment of fungal infections of the skin and nails including candidiasis, tinea, and pityriasis versicolor.

Pharmacodynamics & Pharmacokinetics

Miconazole is an antifungal agent which also has antibacterial activity against some Gram-positive bacteria. It acts by damaging fungal cell membrane and altering its permeability resulting in loss of essential intracellular elements. It also inhibits biosynthesis of triglycerides and phospholipids by fungi and oxidative and peroxidative enzyme activity, resulting in deterioration of subcellular organelles and cellular necrosis.

It is absorbed to a small extent percutaneously.

Adverse effects

Local irritation and occasionally sensitisation may occur at the site of application.

Precautions

- Avoid contact with the eyes.
- Do not apply occlusive dressing over this medication unless directed to do so by physician.
- Do not use in children under 2 years of age except under the advice and supervision of a doctor.
- If irritation occurs or if there is no improvement within 4 weeks, discontinue use and consult a doctor or pharmacist.
- In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see Interactions).

Interactions

Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and the anticoagulant effect should be monitored (see Warnings and Precautions).

Dosage and Administration

Skin infections — Apply sparingly and gently rub into the affected area once or twice daily until lesion has completely disappeared.

Nail infections — Apply sparingly to the affected nail twice daily; combine with oral antimycotic or with a non-porous dressing, if necessary.

Medication to be continued for full course of treatment.

Note: To reduce the possibility of recurrence, fungal infections should be treated for 2 weeks.

Storage

: Keep container tightly closed. Keep out of reach of children.
Store in a cool, dry place below 30°C. Protect from light.

Presentation / Packing

: Cream 2% x 15gm.

Product registration holder & Manufacturer

: **PRIME PHARMACEUTICAL SDN. BHD.**
1505, Lorong Perusahaan Utama 1,
Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

11/11/2022