

SAME SIZE ARTWORK
LEAFLET SIZE: 140 mm x 88 mm

Symptoms and Treatment for Overdosage and Antidote (s) Acute overdose with topical application of Clotrimazole is unlikely and it is not expected to represent a life-threatening situation.		<i>For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory</i>	
Packing / Pack Sizes Tube of 20 g, 60 g.		candid * Cream	
		SPACE FOR PHARMACODE	
Shelf Life 4 years		Composition Clotrimazole USP 1% w/w In water miscible base q.s.	
Storage Conditions, User Instructions and Pharmaceutical Precautions Store below 30°Protect from light. MAL19900296XZ		Therapeutic Class D01AC01	
Date of revision: 29/12/2020		Description A printed carton with an insert and a printed aluminium collapsible tube containing a white semi-solid cream.	
Manufactured by : glenmark PHARMACEUTICALS LTD. Plot No. E-37, 39, D-Road, MIDC, Satpur, Nashik - 422 007, Maharashtra State, India. © Trade Mark		Product Registration Holder: GLENMARK PHARMACEUTICALS (Malaysia) Sdn Bhd (660397-W) D-31-02, Menara Suezcap 1, No. 2, Jalan Kerinchi, 59200 Kuala Lumpur, Malaysia	

SAME SIZE ARTWORK
LEAFLET SIZE: 140 mm x 88 mm

(Pityrosporum orbiculare), Paracoccidioides brasiliensis, Sporothrix schenckii, Pseudallescheria, Medurella and Phialophora spp. Clotrimazole also has antibacterial action and some strains of Staphylococcus aureus and Streptococcus pyogenes have been reported to be sensitive. Clotrimazole is believed to exert its activity on the cell membrane of the fungus. It appears that the drug inhibits protein synthesis and so influences the utilisation and the incorporation of the acylated amino acids essential for fungi. Pharmacokinetics Clotrimazole appears to be well absorbed in humans following oral administration and is eliminated mainly as inactive metabolites. Following topical and vaginal administration, however, Clotrimazole appears to be minimally absorbed. Six hours after the application of radioactive Clotrimazole 1% cream and 1% solution onto intact and acutely inflamed skin, the concentration of Clotrimazole varied from 100 mcg / cm³ in the stratum corneum to 0.5 to 1 mcg / cm³ in the stratum reticulare, and 0.1 mcg / cm³ in the subcutis. No measurable amount of radioactivity (≤ 0.001 mcg/ml) was found in the serum within 48 hours after application under occlusive dressing of 0.5 ml of the solution or 0.8 g of the cream. Only 0.5% or less of the applied radioactivity was excreted in the urine. Indications Indicated for the treatment of : ● Tinea Corporis ● Tinea Cruris ● Tinea Pedis ● Tinea Versicolor ● Cutaneous Candidiasis Contraindications Except for possible hypersensitivity reactions, there are no contraindications.	Side Effects / Adverse Reactions The local tolerance of CANDID Cream is excellent and no systemic effects are seen with topical use. Rarely patients may experience local irritation. This generally disappears on continuing the therapy. Drug Interactions None reported. Precautions / Warnings For irritation or sensitivity caused with the use of Clotrimazole; treatment should be discontinued and appropriate therapy instituted. The patient should be advised to use the medication for the full treatment time even though the symptoms may have improved. This medicine should be used in the first trimester of pregnancy after consideration of the benefit/risk relation and by physician's decision. It is not known whether Clotrimazole is excreted in human being's milk. As many drugs are excreted in human being's milk, caution should be exercised with Clotrimazole used by nursing mothers. Pregnancy and Lactation This medicine should be used in the first trimester of pregnancy after consideration of the benefit / risk relation and by physician's decision. It is not known whether Clotrimazole is excreted in human being's milk. As many drugs are excreted in human being's milk, caution should be exercised with Clotrimazole used by nursing mothers. Recommended Dosage, Dosage Schedule and Route of Administration The affected area should be washed and thoroughly dried. CANDID Cream is evenly rubbed into the affected area 2-3 times daily. Treatment should continue for 2-4 weeks after the disappearance of all signs of infection to ensure mycological cure and prevent relapse.
---	--

ICONGRAPHICS CODE:

PANTONE SHADE
 PANTONE BLACK PROCESS C
Supersedes Artwork Code : PE56445

PHARMACODE:

PE00000 MY