

33 mm

25 mm

INSTRUCTIONS FOR USE

SPIKE

 Tighten the cap on the nozzle

 The spike in the cap will pierce the tip of the vial.

 SPIKE

Tighten the cap on the nozzle

The spike in the cap will pierce the tip of the valve



Dispense drops with gentle pressure.
Close the cap properly after every use.



**BROAD SPECTRUM ANTIMYCOTIC
WITH FUNGICIDAL ACTION**

Clotrimazole USP	1% w/v
Non-aqueous Base	q.s.

D01AC01

Description
A printed carton with an insert and a labeled, LDPE bottle fitted with a plug and cream coloured cap containing a clear, colourless, viscous liquid.

Clotrimazole has broad spectrum fungicidal activity against Dermatophytes (Trichophyton, Epidermophyton and Microsporum spp.), Candida and aspergillus spp., Malassezia furfur and other fungi. It also inhibits some strains of Gram-positive and Gram-negative bacteria.

Pharmacokinetics, Pharmacodynamics

Clotrimazole is an imidazole antifungal agent and is effective against a wide variety of fungi including Epidermophyton, Microsporum and Trichophyton spp., Coccidioides immitis, Histoplasma capsulatum, Aspergillus and Candida spp. Other fungi responding to Clotrimazole include Cryptococcus neoformans, Malassezia furfur (Pityrosporum orbiculare), Paracoccidioides brasiliensis, Sporothrix schenckii, Psuedallescheria, Medurella and Phialophora spp. Clotrimazole also has antibacterial action and some strains of Staphylococcus aureus

Product Registration Holder:
Glenmark Pharmaceuticals
(Malaysia) Sdn Bhd (660397-W)
D-31-02, Menara Suezcap 1,
No. 2, Jalan Kerinchi,
59200 Kuala Lumpur, Malaysia

Glenmark PHARMACEUTICALS LTD.
Plot No. E-37, 39, D-Road, MIDC, Satpur,
Nashik - 422 007, Maharashtra State, India
Trade Mark

PE00000 MY

and *Streptococcus pyogenes* have. Clotrimazole is believed to exert its antifungal effect by blocking the ergosterol synthesis in the cell membrane of the fungus. It appears that the drug also influences the utilisation and the interconversions of the sterol precursors essential for fungi.

Pharmacokinetics

Clotrimazole appears to be well absorbed after oral administration and is eliminated mainly in the urine. Following topical and vaginal administration, the drug appears to be minimally absorbed. After application of a 1% cream of radioactive Clotrimazole to an acutely inflamed skin, the concentration of the drug in the stratum corneum was $100 \text{ mcg} / \text{cm}^2$ in the stratum corneum, and $0.1 \text{ mcg} / \text{cm}^2$ in the stratum reticulare, and $0.001 \text{ mcg} / \text{cm}^2$ in the stratum granulosum. The amount of radioactivity ($\leq 0.001 \text{ mcg}$) excreted in the urine within 48 hours after application was 0.001 mcg . After application of the solution or 0.8 g of the cream, the radioactivity was excreted in the u-

and *Streptococcus pyogenes* have been reported to be sensitive. Clotrimazole is believed to exert its activity on the cell membrane of the fungus. It appears that the drug inhibits protein synthesis and so influences the utilisation and the incorporation of the acylated amino acids essential for fungi.

Pharmacokinetics
Clotrimazole appears to be well absorbed in humans following oral administration and is eliminated mainly as inactive metabolites. Following topical and vaginal administration, however, Clotrimazole appears to be minimally absorbed. Six hours after the application of radioactive Clotrimazole 1% cream and 1% solution onto intact and acutely inflamed skin, the concentration of Clotrimazole varied from 100 mcg / cm² in the stratum corneum to 0.5 to 1 mcg / cm² in the stratum reticulare, and 0.1 mcg / cm² in the subcutis. No measurable amount of radioactivity (≤ 0.001 mcg / ml) was found in the serum within 48 hours after application under occlusive dressing of 0.5 ml of the solution or 0.8 g of the cream. Only 0.5% or less of the applied radioactivity was excreted in the urine.

CANDID Lotion can be used, with benefit in the following conditions:

- Tinea Capitis
- Tinea Barbae
- Tinea Unguium
- Tinea Corporis
- Tinea Cruris
- Tinea Pedis
- Tinea Versicolor
- Candidal Intertrigo
- Candidal Paronychia
- Candidal Vulvitis
- Candidal Balanitis
- Anogenital Candidiasis
- Fungal Diaper Dermatitis
- Erythrasma

CANDID Lotion is ideal for the treatment of large lesions, hairy and intertriginous skin areas and for the nails.

Contraindications
Except for rare cases of hypersensitivity, there are no contraindications.

The local tolerance of **CANDID Lotion** is excellent and no systemic effects are seen with topical use. Rarely patients may experience local irritation. This generally disappears on continuing the therapy.

Drug interactions
None reported

Precautions, warnings
For irritation or sensitivity caused with the use of Clotrimazole; treatment should be discontinued and appropriate therapy instituted.

The patient should be advised to use the medication for the full treatment time even though the symptoms may have improved. This medicine should be used in the first trimester of pregnancy after consideration of the benefit/risk relation and by physician's decision. It is not known whether Clotrimazole is excreted in human being's milk. As many drugs are excreted in human being's milk, caution should be exercised with Clotrimazole used by nursing mothers.

This medicine should be used in the first trimester of pregnancy after consideration of the benefit / risk relation and by physician's decision. It is not known whether Clotrimazole is excreted in human being's milk. As many drugs are excreted in human being's milk, caution should be exercised with Clotrimazole used by nursing mothers.

The affected area should be washed and thoroughly dried. **CANDID Lotion** is evenly rubbed into the affected area 2-3 times daily. Treatment should continue for 2-4 weeks after the disappearance of all signs of infection to ensure mycological cure and prevent relapse.

Acute overdose with topical application of Clotrimazole is unlikely and it is not expected to represent a life-threatening situation.

Bottle of 20 ml.

2 years

Store below 30°C. Protect from light.

MAL19900303XZ

Date of Revision: 30/07/2022

PE00000 MY

1004-1005

PANTONE SHADE

PANTONE
BLACK
PROCESS C

**Supersedes
Artwork Code:
PE50463**

PHARMACODE: