

SAME SIZE ARTWORK
LEAFLET SIZE: 130 mm x 100 mm

33 mm

25 mm

25 mm

Shelf Life
24 months

Storage Conditions, User Instructions and Pharmaceutical Precautions
Store below 30°C. Protect from light
MAL 19900331AZ
Date of Revision: 23/02/2021

SPACE FOR PHARMACODE

INSTRUCTIONS FOR USE

SPIKE
Tighten the cap on the nozzle

The spike in the cap will pierce the tip of the vial.

Dispense drops with gentle pressure.
Close the cap properly after every use.

glenmark
PHARMACEUTICALS LTD.
Plot No. E-37, 39,
D-Road, MIDC, Satpur,
Nashik - 422 007,
Maharashtra State, India.
© Trade Mark

Product Registration Holder
GLENMARK
PHARMACEUTICALS
(Malaysia) Sdn Bhd (660397-W)
D-31-02, Menara Suezcap 1,
No. 2, Jalan Kerinchi,
59200 Kuala Lumpur, Malaysia

MY
PE00000

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

candid*
Ear Drops

Composition
Clotrimazole USP 1% w/v
Propylene Glycol USP Base q.s.

Therapeutic Class
T0130

Description
A printed carton with an insert and a labeled, LDPE bottle fitted with a plug and blue coloured cap containing a clear, colourless, viscous liquid.

Actions and Mode or Mechanism of Action
Clotrimazole has broad spectrum fungicidal activity against Dermatophytes (Trichophyton, Epidermophyton and Microsporum spp.), Candida and Aspergillus spp., Malassezia furfur and other fungi. It also inhibits some strains of Gram-positive and Gram-negative bacteria.

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics)
Pharmacodynamics
Clotrimazole is an imidazole antifungal agent and is effective against a wide variety of fungi including Epidermophyton, Microsporum and Trichophyton spp., Coccidioides immitis, Histoplasma capsulatum, Aspergillus and Candida spp. Other fungi responding to Clotrimazole include Cryptococcus neoformans, Malassezia furfur (Pityrosporum orbiculare), Paracoccidioides brasiliensis, Sporothrix schenckii, Pseudallescheria, Medurella and Phialophora spp.

SAME SIZE ARTWORK
LEAFLET SIZE: 130 mm x 100 mm

Clotrimazole also has antibacterial action and some strains of Staphylococcus aureus and Streptococcus pyogenes have been reported to be sensitive. Clotrimazole is believed to exert its activity on the cell membrane of the fungus. It appears that the drug inhibits protein synthesis and so influences the utilisation and the incorporation of the acylated amino acids essential for fungi.

Pharmacokinetics

Clotrimazole appears to be well absorbed in humans following oral administration and is eliminated mainly as inactive metabolites. Following topical and vaginal administration, however, Clotrimazole appears to be minimally absorbed. Six hours after the application of radioactive Clotrimazole 1% cream and 1% solution onto intact and acutely inflamed skin, the concentration of Clotrimazole varied from 100 mcg / cm² in the stratum corneum to 0.5 to 1 mcg / cm² in the stratum reticulare, and 0.1 mcg / cm² in the subcutis. No measurable amount of radioactivity ($\leq$ 0.001 mcg/ml) was found in the serum within 48 hours after application under occlusive dressing of 0.5 ml of the solution or 0.8 g of the cream. Only 0.5% or less of the applied radioactivity was excreted in the urine.

Indications

- Ootomycosis.
- Concomitant therapy with Antibiotic and Corticosteroid Ear Drops.
- Prophylaxis for swimmers.

Contraindications

Except for possible hypersensitivity reactions, there are no contraindications to CANDID Ear Drops.

Side Effects / Adverse Reactions

The local tolerance of CANDID Ear Drops is excellent and no systemic effects are seen with topical use. Rarely patients may experience local irritation. This generally disappears on continuing the therapy.

Drug Interactions

None reported.

Precautions / Warnings

When the ear drum is perforated, CANDID Ear Drops should be used with caution. This in accordance with the general directions for the use of all ear drops.

This medicine should be used in the first trimester of pregnancy after consideration of the benefit/risk relation and by physician's decision. It is not known whether Clotrimazole is excreted in human being's milk. As many drugs are excreted in human being's milk, caution should be exercised with Clotrimazole used by nursing mothers.

Pregnancy and Lactation

When the ear drum is perforated, CANDID Ear Drops should be used with caution. This in accordance with the general directions for the use of all ear drops. This medicine should be used in the first trimester of pregnancy after consideration of the benefit/risk relation and by physician's decision. It is not known whether Clotrimazole is excreted in human being's milk. As many drugs are excreted human being's milk, caution should be exercised with Clotrimazole used by nursing mothers.

Recommended Dosage, Dosage Schedule and Route of Administration

4-5 drops are instilled in the ear 3-4 times daily or as directed by the physician.

Symptoms and Treatment for Overdosage and Antidote(s)

Acute overdose with topical application of Clotrimazole is unlikely and it is not expected to represent a life-threatening situation.

Packing / Pack Sizes

Bottle of 15 ml.

PE00000 MY

ICONGRAPHICS CODE:

PANTONE SHADE

PANTONE
BLACK
PROCESS C

Supersedes
Artwork Code:
PE19364

PHARMACODE: