

SAME SIZE ARTWORK
LEAFLET SIZE: 88 mm x 186 mm

24 mm

SPACE FOR PHARMACODE

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

candid-v6™

Clotrimazole Vaginal Inserts USP 100 mg

SPACE FOR PHARMACODE

Composition
Each uncoated tablet contains:
Clotrimazole USP 100 mg

Preservatives:
Methyl Paraben USP NF 0.8 mg
Propyl Paraben USP NF 0.2 mg

Therapeutic Class
T0130

Description
A printed carton with an insert, an applicator and a printed aluminium strip containing 6 white, biconvex vaginal, uncoated tablet with monogram on one side and V6 on the other side.

Actions and Mode of Mechanism of Action
Clotrimazole is an imidazole antifungal agent. Clotrimazole exerts its activity on the cell membrane of the fungus. It appears that the drug inhibits protein synthesis and so influences the utilisation and the incorporation of the acylated amino acids essential for fungi.

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics)
Pharmacodynamics
Clotrimazole has a wide spectrum of activity against Dermatophytes (Trichophyton, Microsporum, Epidermophyton), Yeasts (Candida sp., Torulopsis sp.), Mould fungi and other fungi (Aspergillus sp., Malassezia furfur, Nocardia sp., etc.). It also has action against Trichomonas and some strains of Gram - positive bacteria (Streptococci, Staphylococci and Corynebacterium minutissimum) and Gram - negative bacteria (Bacteroides and Haemophilus vaginalis).

Pharmacokinetics
Following topical and vaginal administration, however, Clotrimazole appears to be minimally absorbed. Following intravaginal administration of 100 mg ¹⁴C-Clotrimazole vaginal inserts to nine adult females, an average peak serum level, corresponding to only 0.03 µg equivalents/mL of Clotrimazole, was reached 1 to 2 days after application.

Indications
Vulvovaginal Candidiasis.

Contraindications
Except for possible hypersensitivity reactions, there are no contraindications.

Side Effects / Adverse Reactions
There is virtually no absorption of the active ingredient through the vaginal mucosa. No systemic effects are therefore to be expected. **CANDID-V6** Vaginal Inserts are well tolerated locally.

Drug Interactions
None reported.

Precautions / Warnings
For irritation or sensitivity caused with the use of Clotrimazole; treatment should be discontinued and appropriate therapy instituted. There are no adequate and well-controlled studies in pregnant women during first trimester although in clinical trials, use of vaginally applied Clotrimazole in pregnant women in their second and third trimester has not been associated with ill effects. Clotrimazole should be used during pregnancy

SPACE FOR PHARMACODE

PE00000 MY

SAME SIZE ARTWORK
LEAFLET SIZE: 88 mm x 186 mm

only if clearly needed. It is not known whether Clotrimazole is excreted in human being's milk. As many drugs are excreted in human being's milk, caution should be exercised with Clotrimazole used by nursing mothers. Clotrimazole vaginal inserts may reduce the efficacy and safety of latex contraceptive such as condom & diaphragm.

Recommended Dosage, Dosage Schedule and Route of Administration

Non-Pregnant Patients: Intravaginal, 200 mg., once a day, preferably at bedtime for three consecutive days, or 100 mg., once a day, preferably at bedtime for seven consecutive days.

Pregnant Patients: Intravaginal, 100 mg., once a day, preferably at bedtime for seven consecutive days.

The Insert is to be inserted as deeply as possible into the vagina. This is best achieved with the applicator when lying on the back with legs pulled in a little towards the body.

The treatment should not be carried out during menstruation but should be completed before this begins. For complete relief, it is advisable to apply **CANDID** Cream to the area around the vagina. For the prevention of reinfection, the partner should be treated locally with **CANDID** Cream.

Directions for Use of the Applicator

1. Pull out plunger A until it stops. Place one Vaginal Insert into the applicator B.
2. Insert applicator containing the Insert carefully as deep as possible into the vagina (this is done best by lying on the back).
3. Push plunger A until it stops, thereby depositing the Insert into the vagina. Withdraw the applicator.
4. After use, remove plunger A completely by pulling it out of the applicator B. Thereafter wash both in warm (not boiling) soapy water and dry completely.



Symptoms and Treatment for Overdosage and Antidote(s)

Acute overdose with topical application of Clotrimazole is unlikely and it is not expected to represent a life-threatening situation.

Packing / Pack Sizes

Strip of 6 Inserts with an applicator.
17 Strips of 6 Vaginal Inserts each with 20 applicators.

Shelf Life
4 years

Storage Conditions, User Instructions and Pharmaceutical Precautions
Store below 25°C. Protect from moisture & light.

Keep medicine out of reach of children.

MAL 19900333AZ

Date of Revision: 1 October 2018

Manufactured by
glenmark
PHARMACEUTICALS LTD.
Plot No. E-37, 39, D-Road,
MIDC, Satpur, Nashik - 422 007,
Maharashtra State, India
© Trade Mark

Product Registration Holder:
Glenmark Pharmaceuticals
(Malaysia) Sdn Bhd (660397-W)
D-31-02, Menara Suezcap 1,
No. 2, Jalan Keintih,
59200 Kuala Lumpur, Malaysia

P-EX0000 MY

ICONGRAPHICS 0000

PANTONE SHADE
 PANTONE BLACK PROCESS C
Supersedes Artwork Code:

PHARMACODE: