

SAME SIZE ARTWORK  
LEAFLET SIZE: 88 mm x 186 mm

24 mm

SPACE FOR PHARMCODE

For the use only of a Registered Medical Practitioner  
or a Hospital or a Laboratory

**candid<sup>®</sup>-v 3**  
**Clotrimazole Vaginal Inserts USP 200 mg**

SPACE FOR PHARMCODE

**Composition**  
Each uncoated tablet contains:  
Clotrimazole USP 200 mg

**Preservatives:**  
Methyl Paraben USP NF 0.8 mg  
Propyl Paraben USP NF 0.2 mg

**Therapeutic Class**  
T0130

**Description**  
A printed carton with an insert, an applicator (s) and a printed aluminium strip (s) containing 3 white, biconvex vaginal, uncoated tablets with monogram on one side and V3 on the other side.

**Actions and Mode or Mechanism of Action**  
Clotrimazole is an imidazole antifungal agent. Clotrimazole exerts its activity on the cell membrane of the fungus. It appears that the drug inhibits protein synthesis and so influences the utilisation and the incorporation of the acylated amino acids essential for fungi.

**Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics)**  
**Pharmacodynamics**  
Clotrimazole has a wide spectrum of activity against Dermatophytes (Trichophyton, Microsporum, Epidermophyton), Yeasts (Candida sp., Torulopsis sp.), Mould fungi and other fungi (Aspergillus sp., Malassezia furfur, Nocardia sp., etc.). It also has action against Trichomonas and some strains of Gram - positive bacteria (Streptococci, Staphylococci and Corynebacterium minitissimum) and Gram - negative bacteria (Bacteroides and Haemophilus vaginalis).

**Pharmacokinetics**  
Following topical and vaginal administration, however, Clotrimazole appears to be minimally absorbed. Following intravaginal administration of 100 mg <sup>14</sup>C-Clotrimazole vaginal inserts to nine adult females, an average peak serum level, corresponding to only 0.03 µg equivalents/mL of Clotrimazole, was reached 1 to 2 days after application.

**Indications**  
Vulvovaginal Candidiasis.

**Contraindications**  
Except for possible hypersensitivity reactions, there are no contraindications.

**Side Effects / Adverse Reactions**  
There is virtually no absorption of the active ingredient through the vaginal mucosa. No systemic effects are therefore to be expected. **CANDID-V3** Vaginal Inserts are well tolerated locally.

**Drug Interactions**  
None reported.

**Precautions / Warnings**  
For irritation or sensitivity caused with the use of Clotrimazole; treatment should be discontinued and appropriate therapy instituted. There are no adequate and well-controlled studies in pregnant women during first trimester although in clinical trials, use of vaginally applied Clotrimazole in pregnant women in their second and third trimester has not been associated with ill effects. Clotrimazole should be used during pregnancy only if clearly needed. It is not known whether Clotrimazole is excreted in human being's milk. As many drugs are excreted in human being's milk, caution should be exercised with Clotrimazole used by nursing mothers. Clotrimazole vaginal inserts may reduce the efficacy and safety of latex contraceptive such as condom & diaphragm.


PE56459 MY

SAME SIZE ARTWORK  
LEAFLET SIZE: 88 mm x 186 mm

**Recommended Dosage, Dosage Schedule and Route of Administration**  
**Non-Pregnant Patients :** Intravaginal, 200 mg., once a day, preferably at bedtime for three consecutive days, or 100 mg., once a day, preferably at bedtime for seven consecutive days.  
**Pregnant Patients :** Intravaginal, 100 mg., once a day, preferably at bedtime for seven consecutive days.  
The Insert is to be inserted as deeply as possible into the vagina. This is best achieved with the applicator when lying on the back with legs pulled in a little towards the body.  
The treatment should not be carried out during menstruation but should be completed before this begins. For complete relief, it is advisable to apply **CANDID Cream** to the area around the vagina. For the prevention of reinfection, the partner should be treated locally with **CANDID Cream**.

**Directions for Use of the Applicator**

1. Pull out plunger A until it stops. Place one Vaginal Insert into the applicator B.
2. Insert applicator containing the Insert carefully as deep as possible into the vagina (this is done best by lying on the back).
3. Push plunger A until it stops, thereby depositing the Insert into the vagina. Withdraw the applicator.
4. After use, remove plunger A completely by pulling it out of the applicator B. Thereafter wash both in warm (not boiling) soapy water and dry completely.



**Symptoms and Treatment for Overdosage and Antidote (s)**  
Acute overdose with topical application of Clotrimazole is unlikely and it is not expected to represent a life-threatening situation.

**Packing / Pack Sizes**  
Strip of 3 Inserts with an applicator.  
34 Strips of 3 inserts with 35 applicators.

**Shelf Life**  
4 years

**Storage Conditions, User Instructions and Pharmaceutical Precautions**  
Store below 30°C. Protect from moisture & light.  
Keep medicine out of reach of children.

**Date of Revision :** 23 Jun 2022  
MAL 19900334AZ

Manufactured by:  
**glenmark**  
**PHARMACEUTICALS LTD.**  
Plot no. E-37, 39, D-Road,  
MIDC, Satpur, Nashik - 422 007,  
Maharashtra State, India.  
© Trade Mark

**Product Registration Holder:**  
Glenmark Pharmaceuticals  
(Malaysia) Sdn Bhd (660397-W)  
D-31-02, Menara Suezcap 1,  
No. 2, Jalan Kerinchi,  
59200 Kuala Lumpur, Malaysia

PE00000 MY

ICONGRAPHICS CODE:

**PANTONE SHADE**

  
**PANTONE  
BLACK  
PROCESS C**

**Supersedes  
Artwork Code:**

PHARMACODE: