

24 mm



For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

candid v-1

Clotrimazole Vaginal Inserts USP 500 mg

Composition

Each uncoated tablet contains:
Clotrimazole USP 500 mg

Preservatives:

Methyl Paraben USP NF 0.8 mg
Propyl Paraben USP NF 0.2 mg

Therapeutic Class

T0130

Description

A printed carton with an insert, an applicator (s) and a printed aluminium strip (s) containing 1/6 white, biconvex vaginal, uncoated tablet with monogram on one side.

Actions and Mode of Mechanism of Action

Clotrimazole is an imidazole antifungal agent. Clotrimazole exerts its activity on the cell membrane of the fungus. It appears that the drug inhibits protein synthesis and so influences the utilisation and the incorporation of the acylated amino acids essential for fungi.

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics)

Pharmacodynamics

Clotrimazole has a wide spectrum of activity against Dermatophytes (Trichophyton, Microsporum, Epidermophyton), Yeasts (Candida sp., Torulopsis sp.), Mould fungi and other fungi (Aspergillus sp., Malassezia furfur, Nocardia sp., etc.). It also has action against Trichomonas and some strains of Gram - positive bacteria (Streptococci, Staphylococci and Corynebacterium minitissimum) and Gram - negative bacteria (Bacteroides and Haemophilus vaginalis).

Pharmacokinetics

Following topical and vaginal administration, however, Clotrimazole appears to be minimally absorbed. Following intravaginal administration of 100 mg ¹⁴C-Clotrimazole vaginal tablets to nine adult females, an average peak serum level, corresponding to only 0.03 µg equivalents/mL of Clotrimazole, was reached 1 to 2 days after application.

Indications

Vulvovaginal Candidiasis.

Contraindications

Except for possible hypersensitivity reactions, there are no contraindications.

Side Effects / Adverse Reactions

There is virtually no absorption of the active ingredient through the vaginal mucosa. No systemic effects are therefore to be expected. **CANDID V-1** Vaginal insert are well tolerated locally.

Drug Interactions

None reported.

Precautions / Warnings

For irritation or sensitivity caused with the use of Clotrimazole; treatment should be discontinued and appropriate therapy instituted. Clotrimazole vaginal tablets may reduce the efficacy and safety of latex contraceptive such as condom & diaphragm.

Recommended Dosage, Dosage Schedule and Route of Administration

A single dose treatment is generally adequate. Unless otherwise prescribed by the Doctor, **Candid V-1** Vaginal insert should be inserted preferably on retiring to bed at night.

It is to be inserted as deeply as possible into the vagina. This is best achieved with the applicator when lying on the back with the legs pulled in a little towards the body.

PE65898 MY

The treatment should not be carried out during menstruation but should be completed before this begins. For complete relief, it is advisable to apply **CANDID** Cream to the area around the vagina. For the prevention of reinfection, the partner should be treated locally with **CANDID** Cream.

USE IN SPECIAL POPULATIONS

Pregnancy

Pregnancy Category B

Clotrimazole Vaginal insert have been used safely in patients in their second and third trimesters of pregnancy, but have not been evaluated in patients during their first trimester.

A study of Clotrimazole use in pregnancy did not find an association between congenital abnormalities and use of Clotrimazole (either vaginal pessaries or topical cream) during pregnancy.

Lactation


Excretion of Clotrimazole in breast milk is unknown. Clotrimazole Vaginal insert should be used in lactating women, only if clearly needed.

Paediatric patients

Clotrimazole Vaginal insert should not be used in children under 12 years of age

Directions for Use of the Applicator

1. Pull out plunger A until it stops. Place one Vaginal insert into the applicator B.
2. Insert applicator containing the Tablet carefully as deep as possible into the vagina (this is done best by lying on the back).
3. Push plunger A until it stops, thereby depositing the insert into the vagina. Withdraw the applicator.
4. After use, remove plunger A completely by pulling it out of the applicator B. Thereafter wash both in warm (not boiling) soapy water and dry completely.



Symptoms and Treatment for Overdosage and Antidote (s)

Acute overdose with topical application of Clotrimazole is unlikely and it is not expected to represent a life-threatening situation.

Packing / Pack Sizes

Strip of 1 insert with an applicator.
17 Strips of 6 insert with 20 applicators.

Shelf Life

4 years

Storage Conditions, User Instructions and Pharmaceutical Precautions

Store below 30°C. Protect from moisture & light.

MAL19900335AZ

Revision Date: 31 October 2023

Product Registration Holder:

Glenmark Pharmaceuticals (Malaysia)
Sdn Bhd (660397-W)
D-31-02, Menara Suezcap 1,
No. 2, Jalan Kerinchi,
59200 Kuala Lumpur, Malaysia

Manufactured by:

glenmark
PHARMACEUTICALS LTD.
Plot No. E-37, 39, D-Road,
MIDC, Satpur, Nashik - 422 007,
Maharashtra State, India.
© Trade Mark

MY
PE65898

ICONGRAPHICS CODE: E39075

PANTONE SHADE



PANTONE
BLACK
PROCESS C

**Supersedes
Artwork Code:
PE54854**

PHARMACODE: 898