

DIPRODIN ELIXIR

COMPOSITION:

Each 5ml contains
Diphenhydramine Hydrochloride 12.5mg
Ephedrine Hydrochloride 6mg

Methylparaben 0.15% w/v as preservative.
Also contains Ammonium Chloride 0.32% w/v
and Citric Acid 0.2% w/v

PRESENTATION:

A red colour strawberry flavoured elixir.

INDICATIONS:

For temporary relief of nasal and sinus congestions associated with wet cough, common cold and allergic rhinitis.

PHARMACOLOGY

Mechanism of Action:

Diphenhydramine HCl is an antihistamine and it has anticholinergic properties. Ephedrine HCl is a sympathomimetic agent and has a pronounced stimulating effects on the CNS. It induces peripheral vasoconstriction and caused bronchodilatation. It also has a stimulant action on the respiratory centre.

DOSAGE & ADMINISTRATION:

Adults: 10ml, 4-6 hourly, not exceeding 60ml in 24 hours. Children 6-12 years: 5ml, 4-6 hourly, not exceeding 30ml in 24 hours. Children 2-6 years: 2.5ml, 4-6 hourly, not exceeding 15ml in 24 hours.

CONTRAINDICATION:

Contraindicated in patients sensitive to any of its ingredients.

PRECAUTION:

If medication causes drowsiness, avoid activities requiring alertness. Use with caution in patients with narrow angle glaucoma, pyloroduodenal obstruction, prostatic hypertrophy, hyperthyroidism, cardiovascular disease or hypertension, and in those with a history of bronchial asthma. Antihistamines may enhance the sedative effects of CNS depressants such as barbiturates, hypnotics, narcotic analgesics, alcohol and tranquilisers. Not recommended for children below 2 years old. To be used with caution and upon the advice of a doctor or pharmacist for children between the ages of 2 to 6 years old.

Use in Pregnancy and Lactation:

Consult a physician before taking this medication.

SIDE EFFECTS:

Like other antihistamine, Diphenhydramine HCl may cause sedation, lassitude, dizziness, hypotension, muscular weakness, gastrointestinal disturbances such as nausea, vomiting, diarrhoea or constipation and epigastric pain, headache, blurred vision, tinnitus, elation or depression, irritability, dryness of the mouth, heaviness and weakness of the hands. Side effects due to Ephedrine HCl may include giddiness, sweating, thirst, tachycardia, precordial pain, palpitations, difficulty in micturition, muscular weakness and tremors, anxiety, restlessness and insomnia.

OVERDOSAGE & TREATMENT:

In recent overdosage the stomach should be emptied by aspiration and lavage. Emetics should not be given. Convulsions may be controlled by IV diazepam. Other symptomatic and supportive treatment should be given where necessary.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN JAUHI DARI KANAK-KANAK

PACK QUANTITIES:

Available in 60ml & 100ml PET bottle and 120ml glass bottle.

Further information can be obtained from pharmacist, physician or the manufacturer.

MAL19910073AZ

Product Registration Holder & Manufactured By:



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of Revision: 18/04/23
REGN - 180423