

Pregnancy and Lactation:

Use in pregnancy: Vitamin B preparations have been widely used during pregnancy without apparent ill consequence; animal studies has shown no hazard.

Riboflavine reduces the anti-bacterial activity of solutions of streptomycin, erythromycin tyrothricin, carbomycin and tetracyclines. Metformin reduces intestinal absorption of cyanocobalamin. Impaired absorption of vitamin B12 was also observed with potassium citrate and or calcium gluconate. Pyridoxine reduces the effects of levodopa.

Urine might be discolored due to riboflavine

Side Effects:

No untoward effects occur within vitamin B complex at the recommended dose.

Symptoms and Treatment of Overdose:

No known.

Storage Condition:

Keep container tightly closed. Store in a cool, dry place (below 30°C). Protect from light.

Pack Size:

Plastic bottle containing 30 tablets and 100 tablets.
Blister pack of 1x10's and 100x10's.

Products Registration Number:

MAL19910317XZ

Further information can be obtained from your doctor or pharmacist.

ROYCE®

Product Holder/Manufactured by:

ROYCE PHARMA MFG. SDN.BHD. (650435-X)
PT 1663, Nilai Industrial Estate,
71800 Nilai, Negeri Sembilan, Malaysia.

Revision date: 121017

ROYCE®

BECOMIN TABLET (Vitamins)

Presentation :

Round yellow tablet with breaking line on one side.

Contents:

Each tablet contains:

Thiamine Hydrochloride	10mg
Calcium Panthothenate	1mg
Riboflavin	1mg
Cyanocobalamin	3mcg
Pyridoxine HCl	1mg
Biotin	30mcg
Nicotinamide.....	20mg

Pharmacological Informamtion:

The vitamin B complex comprises a group of water-soluble factors more or less closely associated in their natural occurrence. Nearly every vitamin of the B complex forms part of a coenzyme essential for the metabolism of protein, carbohydrate or fatty acid.

Thiamine is practically devoid of pharmacodynamic actions when given in the therapeutic dose range. Thiamine, when combine with the pyrophosphoric acid in nucleated cells, particularly in the liver, kidneys and white blood cells, is converted in the body to its pyrophosphate (cocarboxylase).

This functions in carbohydrate metabolism as a coenzyme in the decarboxylation of pyruvic and a-ketoglutaric acids and in the utilization of pentose in the hexose monophosphate shunt.

In the presence of thiamine deficiency, pyruvic and lactic acids accumulate in the tissues. The oxidation of a-keto acids is impaired also. The requirement for thiamine is related to metabolic rate and is greatest when carbohydrate is the source of energy. Absorption from the gastrointestinal tract limited,

mainly from the small intestine and the duodenum. Calcium Panthothenate exhibits no outstanding pharmacodynamic actions following its administration to experimental animals or man. The vitamin carries out its function in the body by being converted into coenzyme A, which serves a vital role in metabolism as a coenzyme for a variety of enzyme-catalysed reactions involving transfer of acetyl (2-carbon) groups.

Riboflavin is inactive until phosphorylated; flavin mononucleotide and flavine adenine dinucleotide are its active forms and these are involved as coenzymes in the respiratory chain and in oxidative phosphorylation.

Riboflavin is readily absorbed from the upper gastrointestinal tract. It is distributed in all tissues, but little is stored and tissue concentrations are uniformly low. When riboflavin is ingested in amounts that approximately the minimal daily requirement, only about 9% appears in the urine. The metabolic fate of the remainder is unknown. As the intake of riboflavin is increased above minimal requirements, larger proportions are excreted unchanged in the urine.

Riboflavin is always present in the feces and probably represent vitamin synthesized by intestinal microorganisms. On low intakes of riboflavin, the amount excreted in the feces is in excess of that ingested. Fecal riboflavin appears to be within the bacterial cells and there is no evidence that it can be absorbed.

The cobalamins are involved in many metabolic systems in man. Cobalamins are essential for normal growth, hematopoiesis, production of all epithelial cells (including those of the gastrointestinal tract) and maintenance of myelin throughout the nervous system. Wherever nucleic acid synthesis occurs, and therefore wherever cells reproduce themselves, cobalamin is required. This requirement is probably proportional to the rapidity of cell proliferation.

The cobalamins are extensively bound to specific plasma proteins, trans cobalamins; trans cobalamin II appears to be involved in the rapid transport of the cobalamins to tissues.

Cobalamins are absorbed from the gastrointestinal tract, but may be irregularly absorbed when given in large therapeutic doses and absorption is impaired in patients with an absence of intrinsic factor, with a malabsorption syndrome or with disease or abnormality of the gut, or after gastrectomy.

Cobalamins are stored in the liver, excreted in the bile and undergo some enterohepatic recycling; part of a dose is excreted in the urine, most of it in the first 8 hours.

Pyridoxine is converted to pyridoxal phosphate, the coenzyme for a variety of metabolic transformations in the body. It is absorbed from the gastrointestinal tract and excreted in the urine as 4-pyridoxic acid.

Biotin is a coenzyme for several enzyme-catalyzed carboxylation reactions during the metabolism of proteins and carbohydrates. Ingested biotin is rapidly absorbed from the gastrointestinal tract and appears in the urine predominantly in the form of free biotin.

Nicotinamide serves a vital role metabolism as coenzymes for a wide variety of proteins involved in tissue respiration. It is rapidly absorbed from the gastrointestinal tract and is widely distributed in the body tissues. The half-life is short.

Indication:

Vitamin B complex deficiency, supplementation of diet in elderly.

Dosage and Administration:

Prophylaxis: One tablet daily.

Treatment: One to two tablets three times daily.

Contraindication:

Known hypersensitivity to any one of the vitamin B complex in the tablet preparation.

Warning and Precautions:

Drug interactions with streptomycin, erythromycin, tetracyclines, metformin, potassium citrate, levodopa etc. Urine may be discolored due to riboflavin.