

Further information can be obtained from your doctor or pharmacist.

ROYCE®

Product Holder/Manufactured by:
ROYCE PHARMA MFG.SDN.BHD. (650435-X)
PT 1663, Nilai Industrial Estate,
71800 Nilai, Negeri Sembilan, Malaysia.

Revision date: 200617

ROYCE®

TRITACID TABLET (Antacids)

Presentation:

A white colour, round, flat face, bevel edge tablet, 12.7mm in diameter, with a breaking line on one side of the tablet and with peppermint flavour.

Content:

Each tablet contains:
Aluminium Hydroxide 120mg
Magnesium Trisilicate 250 mg

Pharmacological Information:

Aluminium Hydroxide reacts with hydrochloride acid in the stomach to form aluminium chloride. This results in the relief from acid pain. Magnesium Trisilicate functions as a nonsystemic antacid.

The gelatinous consistency of the silicon dioxide formed in the stomach is thought to provide an adherent coating that may protect the crater of an ulcer. A fraction of sedimented unreacted excess of the insoluble aluminium hydroxide tends to remain in the stomach to yield a sustained effect.

Magnesium Trisilicate has a slow onset of neutralizing action; in 0.1 N HCl less than 30% reacts in 15 minutes and less than 55% in 30 minutes. Two grams per hour is insufficient to raise the gastric pH above 2.5.

Mucus secretion is stimulated by the irritant action of aluminium ions. Some absorption of aluminium salts occur from the gastrointestinal tract with some excretion in the urine. Some unabsorbed aluminium hydroxide combines with phosphate present in the gut to form insoluble aluminium phosphates and some from carbonates and salts of fatty acids, all these salts are excreted in the faeces.

Only about 5% of magnesium is absorbed and traces of the liberated silica may be absorbed and excreted in the urine.

Indication:

Tritacid is indicated for symptomatic relief of hyperacidity associated with heartburn, aid indigestion, sour stomach and gastric and duodenal ulcers.

Dosage and Administration:

Usual adult dose: Oral, 1 to 2 tablets when required, preferably twenty minutes to one hour after meals and at bedtime.

Usual paediatric dose:

Children up to 6 years of age: Dosage must be individualized by the physician. Children 6 years of age and over, half an adult dose or as directed by the physician.

Contraindication:

Hypophosphataemia

Warning and Precautions:

Do not take this medication:

- if symptoms of appendicitis are present, check with doctor for proper diagnosis.

- If symptoms of inflamed bowel are present

- Within 1 or 32 hours of other oral medication

May interfere with the absorption of tetracyclines.

Sodium content - Alert patients of the possible need for sodium-restricted diet.

Use with caution in patients with renal impairment.

Interactions With Other Medicaments:

Antacids are known, to interface with the absorption of certain drugs, including cimetidine, mexiletine, chlorpromazine, diflunisal, ketoconazole, penicillamine, pivampicillin and the tetracycline.

Pregnancy and Lactation:*Pregnancy*

Aluminium and magnesium containing antacids are considered safe use during the last two trimesters of pregnancy as long as chronic high doses are avoided.

Nursing mothers

Problems in humans have not been documented; although some aluminium and magnesium may be excreted in breast milk, the concentration is not great enough to produce an effect in the neonate.

Side Effects:

With chronic use in renal failure mood or mental changes (neurotoxicity).

With large doses constipation, severe, continuing (fecal impaction), swelling of feet or lower legs.

With prolonged use, bone pain or swelling of wrists or ankles (osteomalacia and osteoporosis due to phosphate depletion).

With prolonged use or large doses, feeling of discomfort, loss, of appetite, muscle weakness or unusual loss of weight (phosphorus depletion syndrome).

With prolonged use or large doses and/ or in renal disease, dizziness or lightheadedness or irregular heartbeat or mood or mental changes or unusual tiredness or weakness (hypermagnesemia or other electrolyte imbalance).

Symptoms & Treatment of Overdose:

Symptoms are unlikely and treatment is rarely required.

Storage Condition:

Keep container tightly closed. Store in a dry place (below 30°C). Protect from light.

Pack Size:

Plastic bottle of 30's.

Product Registration Number:

MAL19910333X