

ROYCE®

Axine Syrup 4mg/5ml

(Bromhexine HCl-Mucolytic)

Presentation:

Yellow coloured, tutti frutti flavoured, clear syrup.

Contents:

Each 5ml contains :

Bromhexine Hydrochloride 4mg

Preservatives:

Methyl paraben 0.1% w/v

Propyl paraben 0.01% w/v

Pharmacological Information:

Bromhexine Hydrochloride has been reported to change the structure of bronchial secretions and to increase the volume and reduce the viscosity of sputum.

It is well absorbed after oral administration. The peak plasma concentration is reached within about an hour. It is also rapidly distributed. It acts through the sulphate and glucuronic acid conjugation.

After an oral dose, about 70% is excreted within 24 hours and in 5 days, about 90% is excreted in the urine and 4% in the faeces. After an intravenous dose, about 50% is excreted in urine in 24 hours and in 5 days, about 80% is excreted in the urine and 5% in faeces.

For further information on absorption, distribution and excretion refer to R. Jauch and R. Hankwitz Arzneimittel Forsch.

Indication:

Reduction of sputum viscosity.

Dosage and Administration:

Adults: Two 5 ml spoonful measure three times daily, not exceeding 48 mg/day.

Children (6 to 12 years): 5 ml three times daily.

Contraindication:

In patients hypersensitive to Bromhexine Hydrochloride.

Warning and Precautions:

Should be administered with caution to patients with gastric ulceration.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

Interaction With Other Medicaments:

Bromhexine has been shown to increase oxytetracycline concentration in sputum and in nasal mucus.

Pregnancy and Lactation:

No report has been reported.

Side Effects:

Gastro-intestinal side effects may occur occasionally and a transient rise in serum aminotransferase values has been reported.

Immune System Disorders

Frequency not known: Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorders

Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

Symptoms and Treatment of Overdose:

See Side Effects.

Storage Condition:

Store in a dry place (below 30°C). Keep container tightly closed. Protect from light.

Pack Size:

Plastic bottles containing 60ml, 90ml, 100ml and 120ml.

Product Registration Number:

MAL19910346XZ

Further information can be obtained from your doctor or pharmacist.

**Product Holder / Manufactured by:**

ROYCE PHARMA MFG. SDN.BHD. (650435-X)

PT 1663, Nilai Industrial Estate,

71800 Nilai, Negeri Sembilan, Malaysia.

Revision date: 211019