

DYSOLVON TABLET 8MG

MAL19910628XZ

DESCRIPTION:

TABLET

Colour : Yellow
Shape : Round/Flat/Scored
Coating: Uncoated

EACH TABLET CONTAINS:

Bromhexine Hydrochloride8 mg

PHARMACODYNAMICS:

Bromhexine Hydrochloride has been reported to change the structure of bronchial secretions and to increase the volume and reduce the viscosity of sputum. It is administered in bronchitis and other respiratory conditions as an aid to expectoration, with variable.

PHARMACOKINETICS:

Bromhexine Hydrochloride is absorbed from the gastrointestinal tract and peak plasma concentrations have occurred after 1 hour. Most of a dose is excreted in the urine mainly as metabolites; only a small amount is excreted in the faeces. Ambroxol is a metabolite of Bromhexine.

INDICATIONS:

Administered in bronchitis and other respiratory conditions to aid expectoration.

RECOMMENDED DOSAGE:

Adults: 2 tablets, 3 to 4 times daily.
Children below 5 years: ½ tablet twice daily.
Children 5 to 10 years: 1 tablet twice daily.

ROUTE OF ADMINISTRATION:

FOR ORAL USE ONLY

CONTRAINDICATION:

Bromhexine should not be used in patients known to be hypersensitive to Bromhexine or other components of the product

PRECAUTIONS/WARNINGS:

Bromhexine should be given cautiously to patient with gastric ulceration.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

DRUG INTERACTIONS:

Administration of Bromhexine together with antibiotics (Amoxicillin, Cefuroxime, Erythromycin, Doxycycline) leads to higher concentrations of the antibiotics in the lungs.

PREGNANCY AND LACTATION:

Available data finds no ill effects following oral administration. Nonetheless, the usual precautions regarding medication use during pregnancy should be practised.

Bromhexine is expected to enter breast milk, and therefore should be avoided during lactation.

SIDE EFFECTS/ADVERSE REACTIONS:

Gastrointestinal side effects may occur occasionally a transient rise in serum transaminase values has been reported.

Immune System Disorders

Frequency not known: Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorders

Frequency not known: Severe skin reactions (including Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP)).

PACKING/PACK SIZE(S):

Blister-pack of 2x10's, 10x10's, 20x10's and 100x10's.

**JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN**

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. (65683-V)
2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, MALAYSIA

PI1NT D030-05
Latest Revision: Nov 2021