

DESCRIPTION:

Chewable Tablet

Colour : White

Shape : Flat, Round, Scored With Logo

Flavour : Peppermint

**EACH TABLET CONTAINS:**

Aluminium Hydroxide.....250 mg

Magnesium Hydroxide250 mg

Simethicone.....50 mg

PHARMACODYNAMICS:

Aluminium is a non systemic antacid; it reacts chemically to neutralise or buffer existing quantities of stomach acid. Simethicone's antifoam properties may enhance the effectiveness of antacid in treating gastric disturbances.

PHARMACOKINETICS:

The insoluble Aluminium compounds that constitute Aluminium Hydroxide mixture are slowly but perhaps incompletely converted to Aluminium Chloride in the stomach. Some absorption of soluble Aluminium salts occurs from the gastrointestinal tract with some excretion in the urine. Some unabsorbed Aluminium Hydroxide combines with phosphates present in the gut to form insoluble Aluminium Phosphates and some form carbonates and salts of fatty acid; all these salts are excreted in the faeces.

Magnesium Chloride and hydrated silica gel are formed during neutralisation. About 5% of the Magnesium are absorbed and traces of the liberated silica may be absorbed and excreted in the urine.

Simethicone is excreted in the faeces as uncharged drug.

INDICATIONS:

Symptomatic relief of peptic ulcer, gastric hyperacidity, dyspepsia, and as an antifatulent.

RECOMMENDED DOSE:

Adults : 2 tablets.

Children 7 to 12 years : 1 tablet.

All to be taken 3 to 4 times daily as required.

ROUTE OF ADMINISTRATION:

FOR ORAL USE ONLY

CONTRAINDICATIONS:

Antacids should not be given to young children (up to 6 years of age) unless prescribed by a physician.

The medication should not be used in severe renal function impairment, appendicitis or intestinal obstruction.

WARNINGS AND PRECAUTIONS:

The medication may interfere with or reduces the absorption of a number of other drugs including anticholinergic agents, barbiturates, digoxin, quinine, warfarin, tetracyclines and vitamins.

The effects of Aluminium Hydroxide on phosphate should always be carefully considered.

SIDE EFFECTS:

Aluminium Hydroxide is astringent and may cause, vomiting and constipation. Increases in plasma-concentration of Aluminium are toxic hazard of Aluminium Hydroxide given to patients with poor renal function, or to those on dialysis, and encephalopathy and dementia have occurred.

SYMPTOMS AND TREATMENT OF OVERDOSE:

The adverse effects of systemic Magnesium poisoning are treated by the intravenous administration of Calcium salts. Dialysis and correction of fluid deficit may be of value. Other supportive measures may be required to normalise the electrolyte imbalance as a result of ingestion of large quantities of the medication.

PACKING/PACK SIZE(S):

Plastic bottles of 20's and 200's.

INDIKASI:

Untuk mengurangkan simptom-simptom ulser peptik, asid yang berlebihan, ketidakhadaman dan sebagai antifatulent.

DOS:

Orange dewasa : 2 tablet.

Kanak-kanak 7 - 12 tahun : 1 tablet.

Diambil 3 - 4 kali sehari.

JAUIH UBAT DARIPADA KANAK-KANAK**KEEP OUT OF REACH OF CHILDREN**

功能 : 缓解消化性溃疡, 胃酸过多, 消化不良的症状, 和胃部气胀。

服法 : 成人 : 每次 2 片。
儿童 7 至 12 岁 : 每次 1 片。
每日服用 3 至 4 次。

MANUFACTURER/PRODUCT REGISTRATION HOLDER:**DYNAPHARM (M) SDN. BHD.** 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, Malaysia.

PI1NT D019-03

Latest Revision: Jul 2022