

DYNA U SUSPENSION

MAL19910648X

DESCRIPTION:

SUSPENSION

Colour : White

Flavour : Peppermint

CONTENT:

EACH 5 ML CONTAINS:

Aluminium Hydroxide 200 mg

Magnesium Hydroxide 200 mg

Simethicone 30 mg

PHARMACODYNAMICS:

Aluminium and Magnesium Hydroxide react chemically to neutralize or buffer existing quantities of stomach acid but it has no direct on its output. This action results in increased pH value of the stomach contents. Between pH 7 and 8, pepsin is irreversibly inactivated.

Simethicone has antifoaming properties. It changes the surface tension of gas bubbles causing them to coalesce.

PHARMACOKINETICS:

Small amounts of the Aluminium in Aluminium Hydroxide are absorbed from the intestine. Approximately 10% of the Magnesium in Magnesium Hydroxide is absorbed from the intestine. Excretion is renal and fecal. 15 to 39% of the salts formed are absorbed and excreted by the kidneys. The excretion of Simethicone is fecal, as the unchanged drug.

INDICATION:

Gastritis, hyperacidity and flatulence.

RECOMMENDED DOSE:

Adults: 1 to 2 teaspoonfuls (5 ml to 10 ml). To be taken after meals and at bedtime or when required.

ROUTE OF ADMINISTRATION:

FOR ORAL USE ONLY

CONTRAINDICATION:

This preparation should not be used in cases of renal impairment, hypophosphatemia, appendicitis or symptoms, undiagnosed gastrointestinal or rectal bleeding, chronic diarrhoea and intestinal obstruction.

WARNING AND PRECAUTIONS:

This preparation may alter the absorption of other drugs and therefore drugs such as tetracyclines or iron salts should be given 1 to 2 hours apart. The maximum dosage should not be taken for more than 2 weeks except under the advice or supervision of a physician. Overuse may inhibit the enzymatic activity of pepsin which is ineffective at pH 4 and above.

This preparation contains Sodium Metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.

SIDE EFFECTS:

Usually mild. Bone pain or swelling of wrists or ankles with prolonged use, unusual tiredness or weakness due to hypermagnesemia or electrolyte imbalance, constipation or diarrhoea, occasional nausea or vomiting.

PACKING/PACK SIZE(S):

Plastic bottle of 120 ml.

JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN
SHAKE WELL BEFORE USE

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, Malaysia.

PI1NL D003-03

Latest Revision: Oct 2024