

DEXCHLORMINE SYRUP

MAL19910823AZ

DESCRIPTION:

SYRUP

Colour : Yellow
Flavour : Banana

CONTENT:

Each 5 ml contains:

Dexchlorpheniramine Maleate 2 mg

Preservatives: Sodium Benzoate 0.1% w/v, Methyl Paraben 0.1% w/v & Propyl Paraben 0.01% w/v.

PHARMACODYNAMICS:

Dexchlorpheniramine Maleate is an antihistamine which acts by competing with histamine for Hi-receptor sites on effector cells. It thereby prevents, but does not reverse responses mediated by histamine alone. The antimuscarinic action provides a drying effect on the nasal mucosa.

PHARMACOKINETICS:

Dexchlorpheniramine Maleate is readily absorbed after oral administration. It is widely distributed in the body. It is metabolised in the liver and excreted in the urine.

INDICATION:

Indicated for the treatment of perennial & seasonal allergic rhinitis; vasomotor rhinitis; allergic conjunctivitis; mild, uncomplicated allergic skin manifestations of urticaria & angioedema; amelioration of allergic reactions to blood or plasma.

RECOMMENDED DOSE:

Adults and children above 12 years of age : 1 teaspoonful (5 ml) every 4 - 6 hours.
Children aged 6 - 11 years : ½ teaspoonful (2.5 ml) every 4 - 6 hours.
Children aged 2 - 5 years : ¼ teaspoonful (1.25 ml) every 4 - 6 hours.

ROUTE OF ADMINISTRATION:

FOR ORAL USE ONLY

CONTRAINDICATION:

- Contraindicated in new-born or premature infants because of the possibility of severe reactions such as convulsions.
- Contraindicated for use in conjunction with MAOI (monoamine oxidase inhibitor) therapy.
- It should not be used to treat lower respiratory tract symptoms.
- Hypersensitivity to the Dexchlorpheniramine Maleate.

WARNING AND PRECAUTIONS:

It may cause drowsiness and dulling of mental alertness. Patients undergoing treatment with these drugs should not take charge of vehicles or machinery. Patients should abstain from alcohol. It may enhance the sedative effect of CNS depressants including alcohol, barbiturates, hypnotics, narcotic analgesics, sedatives and tranquillisers.

It should be used with caution in patients with narrow angle glaucoma, pyloroduodenal obstruction, symptomatic prostatic hypertrophy, bladder neck obstruction, and in patients with a history of bronchial asthma, increased intraocular pressure, hyperthyroidism, cardiovascular disease and hypertension. It should not be used in the third trimester of pregnancy. Antihistamines are more likely to cause dizziness, sedation and hypotension in elderly patients. Overdose of antihistamines may cause hallucinations, convulsions or death especially in infants and children.

DRUG INTERACTION:

It may enhance the sedative effect of CNS depressants including alcohol, barbiturates, hypnotics, narcotic analgesics, sedatives and tranquillisers. Monoamine oxidase inhibitors prolong and intensify the effects of antihistamines. The action of oral anticoagulants may be inhibited by antihistamines.

PREGNANCY AND LACTATION:

Use in pregnancy has not been established. It is not known whether Dexchlorpheniramine is excreted in human milk, therefore caution should be exercised when administered to nursing mothers.

SIDE EFFECTS/ADVERSE REACTIONS:

Slight to moderate drowsiness is the most common side effect. Others include:

- General: Urticaria, drug rash, anaphylactic shock, photosensitivity, excessive perspiration, chills, dryness of mouth, nose and throat.
- Cardiovascular System: Headache, palpitation, tachycardia, hypotension.
- Nervous System: Sedation, dizziness, vertigo, tinnitus, disturbed co-ordination, fatigue, confusion, restlessness, excitation, nervousness, tremor, irritability, insomnia, euphoria, blurred vision and convulsions.
- GIT system: Epigastric distress, anorexia, nausea, vomiting, diarrhoea, constipation.
- Genitourinary system: Urinary frequency, difficult urination, urinary retention, early menses.
- Respiratory system: Thickening of bronchial secretions, tightness of chest, wheezing, nasal stuffiness.

SYMPTOMS AND TREATMENT OF OVERDOSE:

Symptoms of overdose include excitation, convulsions. In severe overdose, the stomach should be emptied by aspiration and lavage. Emetics should not be used. The patient should be kept quiet to minimise the excitation which occurs particularly in children. Convulsions may be controlled with diazepam given intravenously. Other general measures include maintaining respiration with oxygen and assisted respiration provided if necessary.

PACKING/PACK SIZE(S):

Plastic bottle of 60 ml.

**JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN
SHAKE WELL BEFORE USE**

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)
2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, Malaysia.

PI1PL D002-04
Latest Revision: Feb 2022