

DYNAPHAN SYRUP

MAL19910825AZ

DESCRIPTION:

Orange in colour syrupy solution, clear with orange flavour.

EACH 5 ML CONTAINS:

Dextromethorphan HBr 15 mg

PRESERVATIVES: Methyl Paraben 0.1% w/v, Propyl Paraben 0.01% w/v & Sodium Benzoate 0.1% w/v.

PHARMACODYNAMICS:

Dextromethorphan suppresses the cough reflex by a direct central action, probably in the medulla or pons.

PHARMACOKINETICS:

Dextromethorphan HBr is absorbed after oral administration. After an oral dose of 30 mg, 10, 25 and 10% of the dose is detectable in the blood at 1, 2.5, and 6 hours respectively after administration. It is metabolised by N-Demethylation and O-Demethylation followed by Sulphate or Glucuronic Acid conjugation; the major metabolite is (+)-3-hydroxy-N-methylmorphinan and the other metabolites are (+)-3-hydroxymorphinan and possibly (+)-3-methoxymorphinan and their respective conjugates. Up to 56% of a dose is excreted in the urine, about 8% being excreted unchanged in 6 hours, in rats, Dextromethorphan is excreted in the bile and twice as much may be detectable in the faeces as in the urine.

INDICATION:

Symptomatic relief of unproductive cough caused by minor throat and bronchial irritation occurring with colds or inhaled irritants.

RECOMMENDED DOSE:

Adults : 1 to 2 teaspoonfuls (5 ml to 10 ml). Not to exceed 8 teaspoonfuls (40 ml) daily.

Children : 7 - 12 years : ½ to 1 teaspoonfuls (2.5 ml to 5 ml). Not to exceed 4 teaspoonfuls (20 ml) daily.

2 - 6 years : ½ teaspoonfuls (2.5 ml). Not to exceed 2 teaspoonfuls (10 ml) daily.

All to be taken 1 to 4 times daily.

WARNING (When Used For Treatment of Cough and Cold):

1. Not to be used in children less than 2 years of age.
2. To be used with caution and doctor's advice in children 2 to 6 years of age.

CONTRAINDICATION:

It is contraindicated in patients receiving monoamine oxidase inhibitor (MAOI).

WARNING AND PRECAUTIONS:

It should be administered with caution to patients with liver disease and to asthmatic patients. It should not be taken for persistent or chronic coughs or when coughing is accompanied by excessive secretions.

PREGNANCY AND LACTATION:

It is not known whether Dextromethorphan or its metabolites are excreted in breast milk.

There is insufficient information available to determine whether Dextromethorphan has teratogenic potential.

There is no data on the secretion of Dextromethorphan into breast milk and therefore use of this product during lactation should be avoided.

SIDE EFFECTS/ADVERSE REACTIONS:

Drowsiness, dizziness, excitation, mental confusion, gastrointestinal disturbances, nervousness, restlessness or irritability.

SYMPTOMS AND TREATMENT OF OVERDOSE:

Symptoms of overdosage include respiratory depression, ataxia and excitability. Treatment is by naloxone intravenous injection which is an antidote.

PACKING/PACK SIZE(S):

Plastic bottles of 60 ml, 100 ml and 120 ml.

**JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN
SHAKE WELL BEFORE USE**

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, MALAYSIA

PI1PL D023-03
Latest Revision: Jun 2022