

NAME OF THE MEDICINAL PRODUCT

XYLONOR, 150 mg / 1.5 mg per g, oromucosal spray, solution.

QUALITATIVE AND QUANTITATIVE COMPOSITION

1 gram of solution contains 150 mg of Lidocaine and 1.5 mg of Cetrimide.

Excipient(s) with known effect: This medicinal product contains 45.45 g of ethanol 96% per 100 g of solution.

For the full list of excipients, see section on List of Excipients.

One press of the spray delivers 10 mg of lidocaine.

PHARMACEUTICAL FORM

Oromucosal spray, solution.

Clear liquid, not more intensely coloured than slightly brownish-yellow and with a characteristic odour of spearmint.

CLINICAL PARTICULARS

Therapeutic indications

- This medicinal product is indicated for the production of topical anaesthesia in adults, adolescents and children in the buccal cavity, including the following dental procedures:
 - Anaesthetic infiltrations
 - Adjustment and fitting of crowns and bridge pillars
 - juxta- and sub-gingival tartar removal
- Abolition of nausea reflex while making dental impressions.

This medicine is indicated for the adults, the children and the teenagers aged from 6 to 18 years old.

Posology and method of administration

For local use only – Gingival use.

One press of the spray delivers approximately 10 mg of lidocaine base.

Place the spray tip 2 to 4 cm from the area to be anesthetized.

Droplets from one spray press should cover roughly a surface 1cm in diameter. This operation may be repeated at 4 or 5 other oral mucosa sites during the same session.

To decrease the nausea reflex, spray one or more doses of the solution toward the palate.

Do not exceed 20 sprays presses.

Paediatric population

XYLONOR, oromucosal spray, solution is contraindicated for the children under 6 years of age, for the spray use, due to the risk of choking (see Contraindications).

Elderly patients or patients with hepatic function disorders

When liver activity is reduced, the lowest dose leading to efficient anaesthesia shall be used when applied before anaesthetic injection.

Method of administration

For local use only.
Gingival and oral use.

The solution should be applied to a previously dried mucous membrane.

Place the tip of the spray nozzle 2 to 4 cm from the area to be anaesthetized. This operation can be repeated in three different areas of the mouth during the same session. The lowest dose leading to effective anaesthesia should be used.

Saliva aspiration and isolation with a dental cotton roll of the site treated with the local anaesthetic is required. To decrease the nausea reflex, spray one or more doses of the solution toward the palate.

Contraindications

- Hypersensitivity to the active substances, lidocaine and cetrimide, or to any of the excipients listed in section on List of Excipients.
- Hypersensitivity to any local anaesthetics.
- Children (under 6 years of age), because of a risk of choking.

Special warnings and precautions for use

Do not exceed the maximum dose of 1.6 g of solution, equivalent to approximately 20 spray presses during the same anaesthesia ;

Although the systemic passage of lidocaine is expected to be negligible, the medicinal product must be used with caution when applied onto an injured, inflamed or infected area due to the risk of a rapid systemic absorption of lidocaine.

Be sure to limit anaesthesia when spraying the area to be treated.

The accidental orientation of the anaesthetic toward the uvula (soft palate) or the pharynx may trigger a transitory paralysis of these areas and temporary discomfort to the patient. It vanishes quickly.

After application of the anaesthetic to the mucosa, ask the patient to spit any excess product and to avoid swallowing it

Risk of biting trauma (lips, cheeks, tongue) exists but is expected to be very low with the medicinal product due to the limited area of administration. When associated with local injectable anaesthetics, the patient should be told to avoid chewing-gum or eating until sensation is restored.

Tell your patient to inform you if he presents injuries in his mouth because lidocaine may more pass into the blood and interact with other medicines.

This medicinal product contains small amounts of ethanol (alcohol), less than 100 mg per 20 actuations (maximum recommended dose) may be absorbed in the blood.

Interaction with other medicinal products and other forms of interaction

Known interactions which usually occur with lidocaine (beta-blocking agents, inhibitors of CYP1A2, sedatives) are not expected to occur when the product is used locally on the oral mucosa. However, when the oral mucosa is injured, lidocaine may be released into the systemic circulation.

Additive interactions with other local anaesthetics

Toxicity of local anaesthetics is additive. This is not directly applicable considering topical dental anaesthesia but may be a concern when associated with injectable anaesthetics in case of unintended intravascular injection or rapid systemic absorption, especially in children.

The total dose of administered lidocaine should not exceed the maximum recommended dose.

Fertility, pregnancy and lactation

Pregnancy

The medicinal product is applied locally on gingival tissues. No effects during pregnancy are anticipated, since systemic exposure to lidocaine is negligible.

XYLONOR, oromucosal spray, solution can be used during pregnancy.

Breastfeeding

The medicinal product is applied locally on gingival tissues. Lidocaine is excreted in human milk, but at the therapeutic doses used, no effects on the breastfed newborns/infants are anticipated.

XYLONOR, oromucosal spray, solution can be used during breastfeeding.

Fertility

No adverse effects on fertility were observed in preclinical studies.

Effects on ability to drive and use machines

XYLONOR oromucosal spray, solution has no or negligible influence on the ability to drive and use machines.

Undesirable effects

Summary of the safety profile

Adverse reactions following the administration of lidocaine/cetrimide are similar to those observed with other local amide anaesthetics. These adverse reactions are mainly local application site reactions and hypersensitivity reactions.

Systemic adverse reactions are extremely rare with lidocaine ointments. However, they may result from high plasma levels due to excessive dosage, or rapid absorption (see section on Overdose) particularly when associated with local injectable anaesthetics, or may also result from hypersensitivity, idiosyncrasy, or diminished tolerance.

Drowsiness following the administration of lidocaine is usually an early sign of high lidocaine plasma level and may occur as a consequence of rapid absorption.

Serious adverse reactions are generally systemic.

Tabulated list of adverse reactions

The reported adverse reactions come from spontaneous reporting and literature.

The frequency classification follows the convention: Very common ($\geq 1/10$), Common ($\geq 1/100$ to $<1/10$), Uncommon ($\geq 1/1,000$ to $<1/100$), Rare ($\geq 1/10,000$ to $<1/1,000$) and Very rare ($<1/10,000$), “Not known (cannot be estimated from the available data)”.

<i>MedDRA System Organ Class</i>	<i>Frequency</i>	<i>Adverse Reactions</i>
Immune System disorders	Rare	Hypersensitivity including Anaphylactic shock
Nervous System disorders	Not known	Local hypoesthesia
Respiratory, thoracic and mediastinal disorders	Not known	Bronchospasm
Gastrointestinal disorders	Not known	Gingival ulceration

		Oral mucosal exfoliation
Skin and subcutaneous tissue disorders	Not known	Angioedema Erythema Face oedema Rash Pruritus Urticaria
General disorders and administration site conditions	Not known	Application site oedema Application site burn

Overdose

At normal doses and under normal conditions of administration, overdose is unlikely to occur with a product for local use only.

However, caution should be taken when using the product in association with local injectable anaesthetics, as the risk of CNS toxicity and cardiovascular toxicity may occur with high plasma levels of lidocaine due to excessive dosage, or rapid absorption. To date, no cases of overdose have been reported with the product.

Symptomatology:

The following reactions may occur with high plasma levels of lidocaine due to excessive dosage, or rapid absorption in particular when associated with the use of local injectable anaesthetics:

Central Nervous System (CNS):

High plasma concentration may cause CNS stimulation (including seizures) followed by CNS depression (including respiratory arrest) and may be characterized by the following signs and symptoms of escalating severity: circumoral paresthesia, light-headedness, nervousness, anxiety, apprehension, euphoria, confusion, dizziness, drowsiness, hyperacusis, tinnitus, blurred vision, vomiting, nausea, sensations of heat, cold or numbness, twitching, tremors, convulsions, unconsciousness, respiratory depression and arrest. The excitatory manifestations (e.g. twitching, tremors, and convulsions) may be very brief or may not occur at all, in which case the first manifestation of toxicity may be drowsiness merging into unconsciousness and respiratory arrest.

Cardiovascular System:

Cardiovascular manifestations are usually depressant and are characterized by bradycardia, hypotension, arrhythmia and cardiovascular collapse, which may lead to cardiac arrest. Hypertension, tachycardia and angina may be caused by concomitant use with a local injectable anaesthetic containing adrenaline.

Treatment of overdose:

The availability of resuscitation equipment should be ensured before the onset of dental anaesthesia with local anaesthetics.

If signs of acute toxicity are suspected, the medicinal product must immediately be rinsed.

Oxygen should rapidly be administered, if necessary assisted ventilation should be used. Change patient position to supine position if necessary.

In case of cardiac arrest, immediate initiation of cardiopulmonary resuscitation should be performed.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic properties

Pharmacotherapeutic group: Nervous System /Anaesthetics /Local anaesthetics /Amides /Lidocaine combinations. ATC code: N01BB52

The medicinal product is an association of:

- Lidocaine: a local amide anaesthetic. When applied to the oral mucous membrane, it provides surface anaesthesia by controlling painful stimulation occurring in or just beneath the mucosa. The local anaesthetic effect of lidocaine occurs via reversible blockade of nerve fiber impulse propagation.
- Cetrimide: a quaternary ammonium disinfectant with antiseptic properties. This action occurs via protein denaturation, enzyme inactivation and membrane damage of bacteria.

Lidocaine in combination with cetrimide has an onset of action of 2 to 5 minutes and 10 to 20 minutes as duration of action.

Pharmacokinetic properties

Lidocaine:

Absorption:

Results from published studies performed in patients using various topical lidocaine-based preparations applied to healthy oral mucosa show that measured serum lidocaine levels remain well below the toxic range. (< 5 µg/mL). The peak serum level of lidocaine is dose dependent and was found to be between 0.016 µg/mL and 0.35 µg/mL.

Distribution:

Lidocaine is 60% to 80% protein-bound to alpha-1-acid glycoprotein.

Biotransformation:

Lidocaine is principally metabolized in the liver by the cytochrome P450 system. Consequently, a mucosal topical dose of lidocaine is metabolized before entry into the systemic circulation. This accounts for the low plasma lidocaine concentrations following the oral administration of lidocaine.

Excretion:

Lidocaine and its metabolites are excreted by the kidneys, 90% as metabolites and 10% as unchanged drug with a half-life of 100 minutes.

Cetrimide

No pharmacokinetic information regarding cetrimide is available.

Preclinical safety data

The data in animals available are not sufficient to appraise their relevance and do not permit the assessment of the effect of XYLONOR, oromucosal spray, solution on fertility and reproduction and of the genotoxicity and carcinogenicity of the medicinal product.

PHARMACEUTICAL PARTICULARS

List of excipients

Saccharin (E954), spearmint flavour*, dipropylene glycol, ethanol (96 per cent).

*Spearmint flavour composition: mixture of mint and eucalyptus essential oils

Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with any other medicinal products.

Shelf life

36 months.

Special precautions for storage

Do not store above 30°C.

Special precautions for disposal and other handling

Before using the spray, adapt the spray nozzle to the pump and press 3 times vertically to start the pump.

Any unused product or waste material should be disposed of in accordance with local requirements.

PRESENTATION

The drug product is packaged in a metered multidose aerosol, composed of a glass bottle and a pump, on which is fitted a diffuser nozzle. When the diffuser nozzle is not fitted to the pump, the bottle is capped with a plastic cap, which is not in contact with the solution

PRODUCT REGISTRATION HOLDER

N.K. LUCK SDN BHD
91A JALAN PUDU LAMA
50200 KUALA LUMPUR
MALAYSIA

MANUFACTURER

SEPTODONT
58, RUE DU PONT DE CRETEIL
94100 SAINT-MAUR DES FOSSES

Date of Revision

March 2022