



NEUROBION® coated tablets

81641.K029

Vitamins B₁, B₆, and B₁₂

Active ingredients: Vitamins of the B group (B₁, B₆, B₁₂)

Composition

Each coated tablet contains, as active ingredients, 100 mg vitamin B₁ (thiamine disulfide), 200 mg vitamin B₆ (pyridoxine hydrochloride), 200 microgram vitamin B₁₂ (cyanocobalamin).

Excipients: montan-glycol wax, acacia, povidone, methyl cellulose, magnesium stearate, purified water, calcium carbonate, glycerol, mannitol, colloidal anhydrous silica, kaolin, gelatin, titanium dioxide (E 171), sodium starch glycolate, talc, sucrose

Product description

Almost white shiny round coated tablets, highly convex on both sides.

Properties

NEUROBION® is a combination of three essential neurotropic vitamins (B₁, B₆ and B₁₂) in high dosage. Vitamins B₁, B₆ and B₁₂ are of special importance for the metabolism in the peripheral and central nervous system. Their effect on the regeneration of nerves has been shown in various investigations using the vitamins individually and in combination. Vitamin B₁ plays an important role in major metabolic processes. Vitamin B₆ has an analgesic effect. Vitamin B₁₂ ensures blood cell formation and prevents degenerative processes of the nervous system. Both the individual function and the beneficial biochemical links between the three vitamins justify their combined use.

Indication:

NEUROBION® is indicated for neurological and other disorders associated with disturbance of metabolic functions influenced by B complex vitamins, including diabetic polyneuropathy, alcoholic peripheral neuritis and post influenzal neuropathies.

NEUROBION® is also recommended for the treatment of neuritis and neuralgia of the spinal nerves, especially facial paresis, cervical syndrome, low back pain, ischialgia.

Contraindications

NEUROBION® must not be used in patients hypersensitive to any of the active ingredients or excipients of the product (see 'Composition').

NEUROBION® coated tablets are not suitable for the treatment of children below the age of 14 years due to the high content of active ingredients.

Warnings and precautions

In the literature neuropathies are described under long term intake (6-12 months) of more than 50 mg mean daily dose of vitamin B₆. Therefore, under long-term treatment regular monitoring is recommended.

NEUROBION® coated tablets contain sucrose; therefore its use is not recommended in patients with intolerance to some sugars (i.e. rare hereditary galactose or fructose intolerance, glucose-galactose malabsorption, Lapp lactase deficiency, or sucrose-isomaltase insufficiency).

Pregnancy and lactation

No risks have become known associated with the use of **NEUROBION®** during pregnancy at the recommended dosage.

Vitamins B₁, B₆ and B₁₂ are secreted into human breast milk, but risks of overdose for the infant are not known. In individual cases, high doses of vitamin B₆, i.e. > 600 mg daily, may inhibit the production of breast milk.

Adverse effects

Hypersensitivity reactions to vitamin B₁, such as sweating, tachycardia (rapid heart beat), and skin reactions with itching and urticaria are very rare.

Chromaturia ("reddish urine", appears during the first 8 hours after an administration and typically resolves within 48 hours) may occur.

Gastrointestinal complaints, such as nausea, vomiting, diarrhoea or abdominal pain may occur.

Please speak to your doctor, if such symptoms or other unwanted effects occur.

Interactions

The effect of L-dopa may be reduced when vitamin B₆ is administered concomitantly.

Pyridoxine-antagonists, e.g. isoniazid (INH), cycloserin, penicillamin, hydralazine: the efficacy of vitamin B₆ (pyridoxine) may be decreased.

Loop diuretics, e.g. furosemide: In long-term use, the blood level of thiamine may be reduced.

Long term use of acid lowering agents may lead to Vitamin B₁₂ deficiency.

Dosage and administration

1 coated tablet three times daily to treat moderate cases, or to provide interval and follow-up therapy for a course of injections unless prescribed otherwise by the physician.

NEUROBION® coated tablets are swallowed without chewing with a little liquid with or after meals.

The dosage and the duration of treatment are to be defined by the physician depending on the condition of the patient.

Overdose

Sensorial neuropathy and other sensorial neuropathy syndromes which can be caused by long-term administration of high doses of pyridoxine improve gradually upon vitamin discontinuation.

Effects on ability to drive and use machines

No effects of the product on the ability to drive or use machines are known.

Storage and stability

Do not store above 25°C.

Do not use after the expiry date. Keep in a dry place.

Keep medicines out of the reach of children.

Presentations

10, 30, 60, 90, 100 and 500 coated tablets.

Not all containers or pack sizes may be registered or marketed.

Date of revision of text

June 2019 (based on CCDS version 3.0)

Name & address of manufacturer

P&G Health Austria GmbH & Co. OG,
Spittal, Austria.