

HOSOLVON TABLET

VIHOS20-2var (MY)

DESCRIPTION

Round, 7.0 mm in diameter, white to off-white uncoated tablet, shallow convex faces, "HD" embossed and scored on the same face.

COMPOSITION

Each tablet contains:
Bromhexine hydrochloride 8 mg

PHARMACODYNAMICS

Bromhexine hydrochloride is a mucolytic agent which changes the structure of bronchial secretions by rarefaction and fragmentation of the mucopolysaccharide fibres; leading to a reduction in the viscosity of the sputum. It is well absorbed orally and is excreted in the urine mainly as metabolites.

PHARMACOKINETICS

Absorption

Well absorbed after oral administration.

Blood Concentration

Peak plasma concentrations are attained within about 1 hour.

Distribution

Rapidly distributed.

Metabolic Reactions

Sulphate and glucuronic acid conjugation; other routes are not yet elucidated.

Excretion

After an oral dose, about 70% is excreted in the urine in 24 hours and, in 5 days, about 90% is excreted in the urine and 4% in the faeces.

INDICATIONS

Reduction of sputum viscosity.

CONTRAINDICATIONS

There is no absolute contraindication but caution should be observed with gastric ulceration.

WARNINGS AND PRECAUTIONS

- It is advisable not to use this drug during the first trimester of pregnancy.
- Give with caution to patients with gastric ulceration.
- Bromhexine may cause very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP). In most cases, there could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as precaution.

DRUG INTERACTIONS

There is no data on drug interactions/incompatibilities.

MAIN SIDE/ADVERSE EFFECTS

Gastrointestinal disturbances may occur occasionally.

Immune System Disorders:

Anaphylactic reactions including anaphylactic shock.

*Frequency not known

Skin and Subcutaneous Skin Disorders:

Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

*Frequency not known

OVERDOSE AND TREATMENT

Clinical features

Anorexia, nausea, vomiting, abdominal discomfort, diarrhoea, possibly bronchial hypersecretion, hypoventilation and convulsions.

Treat overdose by emesis or gastric lavage, if appropriate; with the necessary symptomatic and supportive measures, if required.

DOSAGE AND ADMINISTRATION

Adults

Orally, 1 to 2 tablets 3 times daily.

Children 6 - 12 years

Orally, 1/2 tablet 3 times daily.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage:

Store below 30°C. Protect from light and moisture.

Presentation/Packing:

Tablet 8 mg x blisters of 3 x 10's and 10 x 10's.

Product Registration Holder: HOVID Bhd.

121, Jalan Tunku Abdul Rahman
(Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Manufactured by: HOVID Bhd.

Lot 56442, 7 1/2 Miles, Jalan Ipoh/Chemor,
31200 Chemor, Perak, Malaysia.

Revision date: August 2020