

HOSOLVON

VIHOS00-0 (MY)

DESCRIPTION

Hosolvon Tablet

Round, 7.0 mm in diameter, white to off-white uncoated tablet, shallow convex faces, "HD" embossed and scored on the same face.

Hosolvon Elixir

Clear, colourless, aniseed and menthol flavoured viscous liquid.

COMPOSITION

Hosolvon Tablet

Each tablet contains: Bromhexine hydrochloride 8 mg

Hosolvon Elixir

Each 5 mL contains: Bromhexine hydrochloride 4 mg

PHARMACODYNAMICS

Bromhexine hydrochloride is a mucolytic agent which changes the structure of bronchial secretions by rarefaction and fragmentation of the mucopolysaccharide fibres; leading to a reduction in the viscosity of the sputum. It is well absorbed orally and is excreted in the urine mainly as metabolites.

PHARMACOKINETICS

Absorption: Well absorbed after oral administration.

Blood Concentration: Peak plasma concentrations are attained within about 1 hour.

Distribution: Rapidly distributed.

Metabolic Reactions: Sulphate and glucuronic acid conjugation; other routes are not yet elucidated.

Excretion: After an oral dose, about 70% is excreted in the urine in 24 hours and, in 5 days, about 90% is excreted in the urine and 4% in the faeces.

INDICATIONS

Reduction of sputum viscosity.

CONTRAINDICATIONS

There is no absolute contraindication but caution should be observed with gastric ulceration.

WARNINGS AND PRECAUTIONS

It is advisable not to use this drug during the first trimester of pregnancy.

Give with caution to patients with gastric ulceration.

DRUG INTERACTIONS

There is no data on drug interactions/incompatibilities.

MAIN SIDE/ADVERSE EFFECTS

Gastrointestinal disturbances may occur occasionally.

OVERDOSE AND TREATMENT

Clinical features

Anorexia, nausea, vomiting, abdominal discomfort, diarrhoea, possibly bronchial hypersecretion, hypoventilation and convulsions.

Treat overdose by emesis or gastric lavage, if appropriate; with the necessary symptomatic and supportive measures, if required.

DOSAGE AND ADMINISTRATION

Hosolvon Tablet

Adults : Orally, 1 to 2 tablets 3 times daily.
Children 6 - 12 years : Orally, 1/2 tablet 3 times daily.

Hosolvon Elixir

Adults : Orally, 10 mL 3 times a day.
Children 6 - 12 years : Orally, 5 mL 3 times a day.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Hosolvon Tablet

Storage:

Store below 30°C. Protect from light and moisture.

Presentation/Packing:

Tablet 8 mg x blisters of 3 x 10's and 10 x 10's.

Product Registration Holder: HOVID Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Malaysia.

Manufactured by: HOVID Bhd.

Lot 56442, 7 1/2 Miles, Jalan Ipoh/Chemor,
31200 Chemor, Perak, Malaysia.

Hosolvon Elixir

Storage:

Store below 30°C. Protect from light and moisture.

Presentation/Packing:

Elixir 4 mg/5 mL x 60 mL, 90 mL, 100 mL and 120 mL.
(Not all presentation available locally)

Product Registration Holder /

Manufactured by: HOVID Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Malaysia.

Revision date: August 2019

hovid