

Front - 70mm

Back - 70mm

REXOM™ BROMHEXINE TABLET 8MG

Name and Strength of Active Ingredient(s)

Each tablet contains:

Bromhexine Hydrochloride BP.....8mg

Product Description

White colour, round with top-score tablet.

Pharmacodynamics

Bromhexine is a mucolytic agent, derived from benzylamine, vasicine and adhatodic acid, alkaloids obtained from the plant *Adhatoda vasica*. It acts by increasing sputum volume and reducing the bronchial secretions in chronic bronchitis patients. It has been reported to induce hydrolytic depolymerization of mucoprotein fibers and to stimulate the activity of ciliated epithelium. Improvements in pulmonary function among bronchitis patients in addition to easier expectoration is one of many significant effect of bromhexine. Most of the dose is excreted in the urine mainly as metabolite, Ambroxol whilst only a small amount is excreted in the faeces.

Pharmacokinetics

Bromhexine HCl is rapidly absorbed from the gastrointestinal tract; peak plasma concentrations occur after about 1 hour. It undergoes extensive first pass metabolism in the liver; its oral bioavailability is stated to be only 20% and widely distributed to body tissues. About 85 to 90% of dose is excreted in the urine mainly as metabolite, Ambroxol. In addition to that, Bromhexine is highly bound to plasma proteins with terminal elimination half-life of 13 to 40 hours. Bromhexine crosses the blood-brain barrier and small amounts crosses the placenta.

Indication

Bromhexine is mucolytic used in the treatment of respiratory disorders associated with productive cough. Generally, it is used in the treatment conditions of the respiratory tract associated with retention of mucoid secretions such as bronchitis and asthma as an aid to expectoration.

Recommended Dosage

Usual adult dose: 8 to 16 mg, 3 times daily

Children 6 to 12 years: 4 mg, 3 times daily

Route of Administration

Oral

Contraindications

Hypersensitivity to bromhexine or other components of the formulation.

Warnings and Precautions

Should be given cautiously to patients with a history of peptic ulcer disease. Should be given cautiously in asthmatic patients. Also, patients with impaired renal and liver function should take this medication cautiously.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration

of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

Interactions with Other Medicaments

Bromhexine may increase the concentration of antibiotics in the lungs; Erythromycin, cephalixin, amoxicillin and thus care is advisable. However, no clinically relevant interactions with other medications have been reported.

Pregnancy & Lactation

Pregnancy: Category A. There is insufficient data on the use of Bromhexine among pregnant women. Thus, the effects are unknown and should only used after seeking proper medical advice from your doctor or pharmacist.

Lactation: Inadequate evidence to determine the infant risk during breastfeeding. However, it is not recommended during the first trimester of pregnancy as studies shows that Bromhexine can penetrate breast milk and thus is not recommended during lactation.

Side Effects

- Gastrointestinal side effects may occur occasionally and a transient rise in serum aminotransferase values has been reported.
- Frequency not known: Immune system disorders like anaphylactic reactions including anaphylactic shock.
- Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).
- Other side effects may include headache, dizziness, sweating.

Symptoms and Treatment of Overdose

None was reported with symptoms of overdose. If they occur, symptomatic treatment should be provided.

Effects on Ability to Drive and Use Machine

Not applicable.

Storage Conditions

Store below 30°C. Protect from light.

Dosage Forms and Packaging

Rexom Bromhexine Tablet 8mg is packed in Alu-PVC strip. Each strip contains 10 tablets. The available packaging for Rexom Bromhexine Tablet 8mg are 3x10's, 10x10's, 50x10's and 100x10's.

Registration Number

MAL19911361XZ

Manufactured by and Product Registration Holder:

KCK PHARMACEUTICAL INDUSTRIES SDN. BHD.

Plot 61, Bayan Lepas Industrial Park Phase 4,

Lintang Bayan Lepas 1, 11900 Bayan Lepas,

Pulau Pinang, Malaysia.

Date of Revision

25 April 2022

P10034-1

115mm