
REXOMTM VITAMIN B-6 TABLET 20MG

Name and Strength of Active Ingredients:

Each tablet contains:

Pyridoxine HCl (B₆)20mg

Product Description

Yellow colour, round with top scored tablet.

Pharmacodynamics

Vitamin B6 (Pyridoxine)

Pyridoxine, once absorbed, is rapidly converted to the co-enzymes pyridoxal phosphate and pyridoxamine phosphate which play an essential role in protein metabolism. Convulsions and hypochromic anaemia have occurred in infants deficient in pyridoxine.

Pharmacokinetics

Vitamin B6 (Pyridoxine)

Pyridoxine is absorbed from the gastro-intestinal tract and converted to the active pyridoxal phosphate which is bound to plasma proteins. It is excreted in the urine as 4-pyridoxic acid.

Indications

Pyridoxine is indicated for adults and children over 12 years old in the treatment of isoniazid-induced peripheral neuritis and pyridoxine deficiency states.

Recommended Dose

Adults and children over 12 years old

Isoniazid neuropathy:-

Prophylaxis 10mg daily

Treatment 50mg three times daily (Adults)

Treatment 30-50mg two to three times a day (children over 12 years)

Deficiency states:-

20-50mg up to three times daily

Elderly

As for adults.

Paediatric population

Little information is available on the use of pyridoxine in children under 12 years. Doses of 4mg/Kg have been used to treat children with familial pyridoxine resistance.

Route of Administration

Oral

Contraindications

Hypersensitivity to the pyridoxine or to any of the excipients.

Warning and Precautions

If symptoms persist or worsen, seek medical advice. Do not exceed the stated dose.

Interactions with Other Medicaments

Pyridoxine antagonises the therapeutic effect of levodopa when it is used without a dopa-decarboxylase inhibitor. Larger doses of pyridoxine can cause a reduction in serum phenytoin and phenobarbitone levels in some patients. Pyridoxine requirements have been reported to be increased by oral contraceptives, hydralazine, isoniazid, cycloserine and penicillamine.

Pregnancy & Lactation*Pregnancy*

There are no or limited amount of data from the use of pyridoxine in pregnant women. Pyridoxine is not recommended during the first trimester of pregnancy.

Breastfeeding

There is insufficient information on the excretion of pyridoxine in human milk. A risk to the breastfed child cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from pyridoxine therapy taking into account the benefit of breast feeding for the child and the benefit of therapy for the woman.

Side Effects

None at recommended dosages. Large daily doses taken over a prolonged period can cause the development of severe peripheral neuropathy.

Symptoms and Treatment of Overdose

Ingestion of 2-3g may cause headache. No treatment necessary.

Storage condition

Store below 30°C. Protect from heat and light. Keep container tightly closed.

Dosage Forms and Packaging Available

REXOM™ Vitamin B-6 Tablet 20mg is packed in Alu-PVDC & Alu-PVC blister pack. Each blister are packed into 10 x 10's and 100 x 10's, in a carton along with a package insert.

REXOM™ Vitamin B-12 Tablet 20mg is also packed in 100's per container.

Registration Number:

MAL19911373X

Manufactured by & Product Registration Holder:**KCK Pharmaceutical Industries Sdn. Bhd.**

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia

Date of Revision:

06 September 2021