
REXOMTM WINA ANTAMAC TABLET

Name and Strength of Active Ingredients:

Each tablet contains:

Aluminium Hydroxide Gel	125mg
Magnesium Trisilicate	250mg
Simethicone	30mg

Product Description

White, round shape with top debossed “ANTAMAC” and bottom “KCK” tablet.

Pharmacodynamics

Antamac tablet is an antacid tablet with slow neutralising action and mild laxative action.

Pharmacokinetics

Magnesium trisilicate is a hydrated magnesium silicate. It is an antacid with general properties similar to those of magnesium hydroxide. When given orally it reacts more slowly with hydrochloric acid in the stomach than magnesium hydroxide. Magnesium trisilicate is often given with aluminium-containing antacids such as aluminium hydroxide, which counteract its laxative effect. Aluminium hydroxide, given orally, slowly reacts with the hydrochloric acid in the stomach to form soluble aluminium chloride, some of which is absorbed. The presence of food or other factors that decrease gastric emptying prolongs the availability of aluminium hydroxide to react and may increase the amount of aluminium chloride formed. Simeticone is a mixture of liquid dimeticones containing finely divided silicon dioxide to enhance the defoaming properties of the silicone. It lowers surface tension and when given by mouth causes bubbles of gas in the gastrointestinal tract to coalesce, thus aiding their dispersion.

Indications

Antamac is used in the symptomatic relief of acid indigestion, dyspepsia, heartburn, gastric pain and flatulence.

Recommended Dose

Adult: 1-2 tablets to be chewed up to 4 times a day before meals or when required

Children over 12 years old: 1-2 tablets to be chewed up to 4 times a day before meals or as directed by doctor.

Route of Administration

Oral

Contraindications

Should not be used in patients who are hypersensitive to any of the active substances or excipients, are severely debilitated or suffering from kidney failure, or hypophosphataemia

Warning and Precautions

Aluminium salts are not, in general, well absorbed from the gastrointestinal tract, and systemic effects are therefore rare in patients with normal renal function. However, care is necessary in patients with chronic renal impairment.

Interactions with Other Medicaments

The absorption of aluminium from the gastrointestinal tract may be enhanced if aluminium compounds are taken with citrates or ascorbic acid. Interactions can be minimised by giving antacids and other medication 2 to 3 hours apart

Pregnancy & Lactation

Problems in pregnancy and lactation in human have not been established. Risk-benefit must be considered. However, aluminium-containing antacids should not be used in neonates and infants because accumulation may lead to increased plasma-aluminium concentrations.

Side Effects

It may cause constipation, diarrhoea and stomach cramps.

Symptoms and Treatment of Overdose

Hypermagnesemia may be corrected by the intravenous injection of 10 to 20ml of 10% Calcium Gluconate injection. Haemodialysis and appropriate correction of fluid and electrolyte balance may be considered.

Effects on Ability to Drive and Use Machine

N/A

Storage Condition

Store below 30°C in a dry place. Protect from direct light.

Dosage Forms and Packaging Available

REXOM™ Wina Antamac Tablet is packed in Alu-PVDC blister pack. Each blister are packed into 10 x 10's and 100 x 10's, in a carton along with a package insert.

REXOM™ Wina Antamac Tablet is also packed in 100's per container.

Date of Revision:

19 July 2025

Registration Number:

MAL19911378X

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia