

Axcel® EVILINE TABLETS

COMPOSITION:

Each tablet contains	
Magnesium Hydroxide	200mg
Aluminium Hydroxide	200mg
Simethicone	20mg

PRESENTATION:

Flat, white, round tablet, 12.9mm in diameter, marked KOTRA logo on one side.

INDICATIONS:

Axcel Eviline Tablet is an antacid / defatulent for symptomatic treatment of peptic ulcer, dyspepsia of functional or organic origin, heartburn in hiatus hernia or pregnancy, flatulence and abdominal distension, gastritis oesophagitis and other conditions where hyperacidity or flatulence may be present.

PHARMACOLOGY:

Mechanism of Actions:

Axcel Eviline Tablet is an antacid used to prevent and relief pain in the symptomatic management of gastric and duodenal ulcers and gastrointestinal reflux by neutralising hydrochloric acid in the gastric secretions. It does not reduce the volume of HCL secreted and may increase it but by increasing the gastric pH which will diminish the activity of pepsin in the gastric secretions. However if the increase of gastric pH is slight, there may be an increase in pepsin activity. Therefore, Axcel Eviline Tablet indirectly suppress peptic activity when given in sufficient quantity to elevate the pH of human gastric contents above 5. Between pH 7 and 8, pepsin is irreversibly inactivated. The presence of antacid increases the volume of gastric juice secreted and the output of HCL. Both aluminium and magnesium hydroxide are classified as non-systemic antacids which neutralise the gastric contents but do not cause systemic alkalosis, because the cationic moiety in the intestine form insoluble basic compounds that are not subsequently absorbed. Both aluminium and magnesium hydroxides react with hydrochloric acid in the stomach to form aluminium chloride and magnesium chloride respectively. The insolubility of both the hydroxide compounds is a desirable property in that fractions of sedimented unreacted excess remain in the stomach to yield a sustained effect. Therefore, the product has a prolonged duration of action. The cathartic effect of magnesium hydroxide is minimized by the aluminium hydroxide, which have constipating effects. Aluminium hydroxide is an effective absorbent and interacts with many drugs in the gastrointestinal tract. Polymethyl siloxane, or simethicone, is an

antifoaming agent with water-repellent properties. It is an adjunct in the treatment of conditions in which gas is a problem, such as flatulence, functional gastric bloating and postoperative gaseous distension.

Pharmacokinetics:

Aluminium hydroxide has a very low equivalent weight and hence a high theoretical neutralising capacity. Proteins, peptides, amino acids and certain dietary organic acids greatly impair the neutralising capacity of Aluminium hydroxide. Although both the hydroxide compounds are classified as non-systemic, some absorption from the gastrointestinal tract does not take place, 5-10% of the magnesium can be absorbed. 17-31% of dietary aluminium is excreted in the urine and a considerable amount is distributed to bone and lung. In renal failure, the administration of aluminium hydroxide can elevate the plasma aluminium concentration. Among the compounds formed in the intestine from aluminium hydroxide are insoluble aluminium phosphates which pass through the intestinal tract unabsorbed, and excreted in the faeces. Ordinarily the absorbed magnesium ion is rapidly excreted by the kidney. The urine may be alkalized, which can decrease the excretion of various weakly basic drugs e.g. quinidine.

DOSAGE & ADMINISTRATION:

For oral administration only.

Two to four tablets, well chewed, four times a day, taken twenty minutes to one hour after meals and at bedtime, or as directed by a physician.

Axcel Eviline Tablet are not recommended for children.

CONTRAINDICATION:

Hypersensitivity to the active substances or to any of the excipients.

- Patients with porphyria.

- Severe renal insufficiency (creatinine clearance less than 30 mL/min).

- Generally contraindicated in children.

- Cachexia.

PRECAUTIONS:

Do not take more than 16 tablets in a 24-hour period or use the maximum dosage for more than two weeks or use if you have kidney disease except under the advice and supervision of a physician.

Aluminium hydroxide may cause constipation and magnesium salts overdose may cause hypomotility of the intestine; large doses of this medicinal product may trigger or aggravate intestinal obstruction and ileus in patients with higher risk, such as those with renal impairment, primary constipation, impaired intestinal motility, infants (0 to 24 months) and the elderly.

Aluminium hydroxide is not well absorbed from the gastrointestinal tract, and systemic effects are therefore rare in patients with normal renal function. However, excessive doses or long-term use, or even normal doses in patients with low-phosphorus diets or in infants (0 to 24 months), may lead to phosphate depletion (due to aluminium-phosphate binding) accompanied by increased bone resorption and hypercalciuria with the risk of osteomalacia. Medical advice is recommended in case of long-term use or in patients at risk of hypophosphatemia.

In patients with renal impairment, plasma levels of aluminium and magnesium tend to increase, which causes hyperalbuminemia and hypermagnesemia respectively.

In these patients, long-term exposure to high doses of aluminium and magnesium salts may lead to encephalopathy, dementia,

microcytic anemia or it may worsen dialysis-induced osteomalacia. Medical monitoring is required in patients with mild to moderate renal failure.

Prolonged use of antacids in patients with mild or moderate renal failure should be avoided.

Administration of this medicinal product is contraindicated in patients with severe forms of renal failure.

Aluminium hydroxide may be unsafe for patients with porphyria who are undergoing hemodialysis.

Pediatric populations

Use of magnesium hydroxide in small children, particularly those with kidney impairment or dehydration, may result in hypermagnesemia.

PREGNANCY AND LACTATION:

There are no available data on use in pregnant women. No conclusion can be drawn regarding whether or not this product is safe for use during pregnancy.

Pregnancy

The medicinal product should only be used if absolutely necessary, under direct medical supervision, after having evaluated the expected benefits for the mother compared with the possible risk to the fetus or infant.

Breast-feeding

Because of the limited maternal absorption when taken in accordance with the indicated dosage regimen, aluminium hydroxide and its magnesium salt combinations are considered compatible with breast-feeding.

SIDE EFFECTS:

The frequency of the undesirable effects listed below is defined using the following conventions:

Common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1000$ to $< 1/100$); rare ($\geq 1/10000$ to $< 1/1000$); very rare ($< 1/10000$); not known (cannot be estimated from the available data).

Immune system disorders:

Frequency not known: angioedema, anaphylactic reactions, hypersensitivity reactions, urticaria, pruritus.

Gastrointestinal disorders:

Uncommon: diarrhea or constipation.

Frequency not known: abdominal pain.

Metabolism and nutrition disorders:

Very rare: hypermagnesemia, which has been observed with prolonged use in patients with kidney impairment.

Frequency not known: hyperalbuminemia, hypophosphatemia, in prolonged use or at high doses, or even normal doses of the medicinal product in patients on low-phosphorus diets, or in infants (0 to 24 months), which may result in increased bone resorption, hypercalciuria or osteomalacia.

DRUG INTERACTION:

Because aluminium and magnesium salts reduce the gastrointestinal absorption of tetracyclines, the use of this product should be avoided during oral tetracycline therapy.

The use of antacids containing aluminium can reduce the absorption of H₂ antagonists, atenolol, cefdinir, cefpodoxime, chloroquine, tetracyclines, diflunisal, digoxin, bisphosphonates, ethambutol,

fluoroquinolones, sodium fluoride, glucocorticoids, indomethacin, isoniazide, kayexalate, ketconazole, levothyroxine, lincosamides, metoprolol, phenothiazine neuroleptics, penicillamines, propranolol, rosuvastatin and iron salts.

-Polystyrene sulfonate (Kayexalate)

Caution is advised when this medicinal product is used concomitantly with polystyrene sulfonate (Kayexalate) due to the potential risk of reduced effectiveness of the resin in binding potassium, of metabolic alkalosis in patients with renal impairment (reported with aluminium hydroxide and magnesium hydroxide) and of intestinal obstruction (reported with aluminium hydroxide).

-Aluminium hydroxide and citrates may cause hyperalbuminemia, especially in patients with renal impairment.

As magnesium hydroxide administration results in urine alkalization, increased excretion of salicylates has been observed in the event of concomitant use.

An interval of at least 2 hours should be allowed (4 for fluoroquinolones), before taking this product to avoid interactions with the other medicinal drugs. Concomitant use with quinidine may increase the serum levels of quinidine and lead to a quinidine overdose.

Concomitant use of aluminium hydroxide and citrates may result in increased aluminium levels, especially in patients with renal impairment. Urine alkalization following magnesium hydroxide administration may affect the excretion of some drugs; increased excretion of salicylates has been observed.

OVERDOSAGE & TREATMENT:

There is very limited experience with deliberate overdose. Cases of overdose with aluminum salts are more likely to occur in patients with severe chronic renal impairment, with the following symptoms: encephalopathy, seizures and dementia, hypermagnesemia. The most frequently reported symptoms of acute overdose with aluminium hydroxide and in combination with magnesium salts include diarrhea, abdominal pain and vomiting. Large doses of this medicinal product may cause or aggravate intestinal obstruction and ileus in patients at risk. Treatment should be symptomatic and include general supportive measures. Aluminium and magnesium are eliminated through the urinary route; treatment of magnesium overdose consists of rehydration and forced diuresis. In case of renal function deficiency, hemodialysis or peritoneal dialysis is necessary.

EFFECTS OF ABILITY TO DRIVE AND USE MACHINE:

Axcel Eviline Tablet has no influence on the ability to drive and use machines.

STORAGE:

Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN JAUHI DARI KANAK-KANAK

PACK QUANTITIES:

Available in blister pack of 10 x 10's and 100 x 10's.

Further information can be obtained from pharmacist, physician or the manufacturer.

Manufacturer and Product Registration Holder :

 **Kotra Pharma (M) Sdn. Bhd.**
No. 1, 2 & 3, Jalan TTC 12, (90082-9)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

MAL19911458X

Date of revision : 26/01/24