



EVILINE FORTE SUSPENSION

COMPOSITION:

Each 5ml contains	
Magnesium Hydroxide	400mg
Aluminium Hydroxide	400mg
Simethicone	40mg

Contains Domiphen Bromide 0.01% w/v as preservative.

PRESENTATION:

A white to off-white coloured suspension with peppermint flavour.

INDICATIONS:

Eviline Forte is an antacid / defatulent for symptomatic treatment of peptic ulcer, dyspepsia of functional or organic origin, heartburn in hiatus hernia or pregnancy, flatulence and abdominal distension, gastension, gastritis oesophagitis and other conditions where hyperacidity or flatulence may be present.

PHARMACOLOGY:

Mechanism of Action:

Eviline Forte is an antacid used to prevent and relief pain in the symptomatic management of gastric and duodenal ulcers and gastro intestinal reflux by neutralising hydrochloric acid in the gastric secretions. It does not reduce the volume of HCl secreted and may increase it but by increasing the gastric pH which will diminish the activity of pepsin in the gastric secretions. However if the increase of gastric pH is slight, there may be an increase in pepsin activity. Therefore, Eviline indirectly suppress peptic activity when given in sufficient quantity to elevate the pH of human gastric contents above 5. Between pH7 and 8, pepsin is irreversibly inactivated. The presence of antacid increases the volume of gastric juice secreted and the output of HCl. Both aluminium and magnesium hydroxide are classified as non-systemic antacids which neutralise the gastric contents but do not cause systemic alkalosis, because the cationic moiety in the intestine form insoluble basic compounds that are not subsequently absorbed. Both aluminium and magnesium hydroxides react with hydrochloric acid in the stomach to form aluminium chloride and magnesium chloride respectively. The insolubility of both the hydroxide compounds is a desirable property

in that fractions of sedimented unreacted excess remain in the stomach to yield a sustained effect. Therefore, the product has a prolonged duration of action. The cathartic effect of magnesium hydroxide is minimized by the aluminium hydroxide, which have constipating effects. Aluminium hydroxide is an effective absorbent and interacts with many drugs in the gastrointestinal tract. Polymethylsiloxane, or simethicone, is an antifoaming agent with water-repellent properties. It is an adjunct in the treatment of conditions in which gas is a problem, such as flatulence, functional gastric bloating and post-operative gaseous distention.

Pharmacokinetics:

Aluminium hydroxide has a very low equivalent weight and hence a high theoretical neutralising capacity. Proteins, peptides, amino acids and certain dietary organic acids greatly impair the neutralising capacity of aluminium hydroxide. Although both the hydroxide compounds are classified as non-systemic, some absorption from the gastrointestinal tract does not take place, 5 - 10% of the magnesium can be absorbed, 17 - 31% of dietary aluminium is excreted in the urine and a considerable amount is distributed to bone and lung. In renal failure, the administration of aluminium hydroxide can elevate the plasma aluminium concentration. Among the compounds formed in the intestine from aluminium hydroxide are insoluble aluminium phosphates which pass through the intestinal tract unabsorbed, and excreted in the feces. Ordinarily the absorbed magnesium ion is rapidly excreted by the kidney. The urine may be alkalized, which can decrease the excretion of various weakly basic drugs e.g. quinidine.

DOSAGE AND ADMINISTRATION :

For oral administration only.
One 5ml spoonful to be taken after meals or whenever symptoms arise.

CONTRAINDICATIONS :

No known contraindications.

PRECAUTIONS :

Care should be taken in patients suffering from hypophosphatemia and impaired renal function. It is probably wise to avoid administration during the first trimester of pregnancy unless there are compelling reasons for its use. Undesirable drug interactions may result, including reduced absorption of tetracycline and vitamins when given concomitantly.

SIDE EFFECTS :

In renal failure, the administration of aluminium hydroxide can elevate the plasma aluminium concentration as high as 53mg/ml. Among compounds form in the intestine from aluminium hydroxide are insoluble aluminium phosphates which pass through the intestinal tract unabsorbed. This can resulting in hypophosphatemia, hypophosphaturia and hypopyrophosphaturia. Phosphate depletion syndrome occasionally observed after chronic aluminium hydroxide ingestion. Patients on the therapeutic regimen for peptic ulcer, usually have high phosphate intake. Therefore ingestion of aluminium hydroxide for this patient results in

decrease phosphate absorption and accompanied by increase in calcium absorption which along with resorption of bone salts causes hypercalciuria. Hypomagnesaemia and hypomagnesiuria occur. Aluminium hydroxide is astringent and may cause nausea, vomiting and constipation. Large doses can cause intestinal obstruction and concretion of fatty acid salts of aluminium in stool. Several cases of intestinal obstruction by a large mass composed of clotted blood has been reported. It also reduces absorption of other drugs including tetracycline and vitamins and could delay the absorption of quinidine. Magnesium hydroxide in some patients has its cathartic effect. Although magnesium hydroxide is classified as a non-systemic antacid, 5 - 10% of magnesium can be absorbed. Retention of any absorbed magnesium can cause neurological, neuromuscular and cardiovascular impairment and even death in persons with renal insufficiency. Ordinarily the absorbed magnesium ion is rapidly excreted by the kidney but the urine may become alkaline. Alkalinization of the urine decrease the excretion of various weakly basic drugs eg. quinidine. Magnesium hydroxide increases the absorption of warfarin.

OVERDOSAGE AND TREATMENT:

Hypophosphataemia may be avoided by concurrent high phosphate intake. The effects of hypermagnesemia may be corrected by the intravenous injection of 10 to 20ml of 10% Calcium Gluconate Injection.

STORAGE :

Keep container well closed. Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN.

JAUHI DARI KANAK-KANAK.

PACK QUANTITIES :

Available in 100ml PET bottle and 120ml amber glass bottle.

Further information can be obtained from pharmacist, physician or the manufacturer.

MAL19911461X

Product Registration Holder & Manufactured By :



Kotra Pharma (M) Sdn. Bhd.
No. 1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

PMSPAXC16105 - 140525(03)
Date of revision: 20/05/25