


BELCID

SUSPENSION
Antacid, Antiflatulent, Demulcent



Composition

BELCID SUSPENSION

Each 15 ml contains :
Aluminum Hydroxide Gel equivalent to
Aluminum Hydroxide 918 mg
Magnesium Hydroxide 300 mg
Simethicone 60 mg

Product Description

Suspension - White with peppermint flavored suspension

Pharmacodynamics

Pharmacological actions:

BELCID has an action of antacid, antiflatulence and demulcent. BELCID acts by neutralizing the gastric acid and also has antifoaming action.

Mechanism of actions:

The clinical use of antacids is based on their ability to increase the pH of gastric secretions: Aluminum hydroxide or oxide reacts with hydrochloric acid to form aluminum chloride and water. Magnesium hydroxide rapidly reacts with hydrochloric acid to form magnesium chloride and water. Simethicone acts by coagulating the mucus-surrounded gas bubbles in the GI tract and expelling them.

Pharmacokinetics

Aluminim hydroxide or oxide is slowly solubilized in the stomach after neutralizing hydrochloric acid to form aluminum chloride, about 17-30% of the aluminum chloride is absorbed and is rapidly excreted by the kidney in patients with normal renal function.

Magnesium hydroxide neutralizes hydrochloric acid in the stomach form magnesium chloride and about 15-30% of the magnesium chloride is absorbed and is rapidly excreted by the kidney in patients with normal renal function. Simethicone is not absorbed from GI tract and is excreted unchanged in feces

Indication

BELCID is used as an antacid, antiflatulence and demulcent in gastric hyperacidity, heartburn and used in the symptomatic relief of peptic and duodenal ulcer pains

Recommended Dose

1 tablespoonful, 4 times daily, 1 hour after meals and at bedtime or as directed by the physician

Route of Administration

Oral

Contraindications

Should not be used in patients with renal failure

Warnings and Precautions

Should be used with caution in patients with renal insufficiency.

Interactions with Other Medicaments

- It may decrease the absorption of tetracycline, digoxin, indomethacin, iron salts when concomitantly administered. Therefore administration of these drugs should be spaced 1-2 hours from doses of BELCID.
- It may increase the absorption of dicumarol which is an anticoagulant

Pregnancy and Lactation

Not known

Side Effects

No adverse effects have been reported

Symptoms and Treatment of Overdose

No symptoms of overdosage have been reported. Should they occur, symptomatic treatment should be provided

Storage Condition

Store in dry place at temperature not exceeding 30°C

Dosage Form and Packaging available

Suspension : 120ml, 240ml



Manufactured by
BIOLAB CO., LTD.
625 Soi 7A Bangpoo Industrial Estate, Sukhumvit Road,
Moo 4, Prakasa, Muang, Samutprakarn 10280, Thailand

Date of Revision : 24.7.2018

EES 019/B

Code : EES019/B

Draft Revision : 0

Final Artwork Revision : B

Name : AW_LEAFLET BELCID NEW MAL.ai

Remark :

Design by

Review by

Approved by

Signature

Signature

Signature

Date

Date

Date

Note :

#0=เพิ่มความ Product description

- ☐ ลูกคำพิจารณาอนุมัติ
- ☐ ทำเสนอราคา
- ☐ ส่งพิมพ์
- ☒ RA ขึ้นทะเบียน
- ☐ อื่นๆ.....

COLOR GUIDE



F-QCP-GE-011

Specification

Size : 120 x 210 MM. (+/- 3 MM.)

Color : พิมพ์ 1 สี คือ สีดำ

Material : ใช้กระดาษปอนด์ น้ำหนัก 55 แกรม

Other :