



PRIMECON TABLET 40mg

Description: White, round biconvex tablet. Diameter = 10.00 mm.

PRIMECON SYRUP 50mg/5ml

Description: Pink coloured, strawberry flavoured syrup.

Contents

Primecon Tablet : Simethicone 40mg/tablet

Primecon Syrup : Simethicone 50mg/5ml

Methylparaben 0.1% w/v

Propylparaben 0.01% w/v

Indications

For treatment of flatulence, abdominal distension and dyspepsia, and other conditions in which the retention of gas may be a problem.

Pharmacology

Simethicone (activated dimethicone/polymethylsiloxane) acts by lowering the surface tension of gas bubbles and causing them to coalesce into a form that can be eliminated more easily by belching or passing flatus. It defoams gastric juice in order to decrease the tendency to gastroesophageal reflux.

Elimination is by faecal excretion.

Adverse effects

None of any clinical significance reported.

Precautions

Safety for use in pregnancy and lactation has not been established.

Dosage and Administration

Adults : Oral, 1 to 2 tablets to be chewed four times a day, after meals and at bedtime, or as needed.

The maximum daily dose should not exceed 480mg.

or

: Oral, 5 to 10ml (1 to 2 teaspoonfuls) four times a day, after meals and at bedtime, or as needed.

The maximum daily dose should not exceed 500mg.

Children : as directed by physician

Storage : Keep container tightly closed.

Store in a cool, dry place below 30°C. Protect from light.

Presentation : Tablet 40mgx100x10's ; Syrup: 50mg/5ml X 60ml

Shelflife : Tablet : 3years

Syrup : 2 years

Product registration : **PRIME PHARMACEUTICAL SDN. BHD.**

holder & Manufacturer : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

21/07/2020