



ALMECON SUSPENSION

Description: White coloured and peppermint flavoured suspension

Content:

Each 5 ml contains: Aluminium Hydroxide 200mg
Magnesium Hydroxide 200mg
Simethicone 20mg
Preservatives: methyl parabens 0.1%w/v
propyl parabens 0.01%w/v

Indications:

Gastric hyperacidity, flatulence and indigestion.

Pharmacology:

Aluminium and magnesium hydroxide, an antacid, reacts chemically to neutralize or buffer existing quantities of stomach acid resulting in increased pH value of stomach contents, thus providing relief of hyperacidity symptoms. Simethicone, an antiflatulent, lowers the surface tension of gas bubbles so that elimination is made easier by belching or passing flatus.

Elimination is by renal and fecal excretions.

Adverse Effects:

Chalky taste or stomach cramps may occur occasionally.

Precautions:

- Caution in patients with renal function impairment, appendicitis, gastrointestinal or rectal bleeding, chronic diarrhea and intestinal obstruction.
- The maximum dosage recommended should not be taken for more than 2 weeks, unless otherwise directed by physician.
- It is safe for use in pregnancy as long as chronic high doses are avoided. However, safety for use in lactation has not been established.

Drug Interactions:

- Concurrent use with oral anticoagulants, antidyskinetics, antimuscarinics, ketoconazole, sucralfate or oral tetracycline may reduce the effectiveness of these medications; patients should be advised to take antacids at least 1 to 2 hours after these medications.
- Caution in patients receiving the following drug therapy. Benzodiazepines, diflunisal, digitalis glycosides, oral loxapine, oral phenothiazines and oral phosphates.

Dosage and administration:

Adults: Oral, 5 – 10ml (1 - 2 teaspoonfuls) to be taken 1 hour after meals and at bedtime.

The maximum daily dose should not exceed 120ml.

Children over 6 years: Oral, 5ml (1 teaspoonful) to be taken 1 hour after meals and at bedtime, or as directed.

Overdosage:

Overuse of antacids may inhibit digestive enzyme activity of pepsin, which is ineffective when the stomach contents reach a pH of 4.0 and above.

Treatment consists of the necessary symptomatic and supportive measures.

Storage : Keep container tightly closed. Keep out of reach of children.
Store in a cool, dry place below 30°C. Protect from light.

Shelflife : 3 years.

Presentation : Suspension x 3.6L (FOR EXPORT ONLY)

Manufactured & : PRIME PHARMACEUTICAL SDN. BHD.

Distributed by : 1505, Lrg. Per. Utama 1, Tmn Perindustrian Bukit Tengah
14000 Bukit Mertajam, Penang.