

VASICAN TABLETS 8 mg

PRODUCT DESCRIPTION:

Green, round and flat tablets, with 8 mm diameter, score-line on one side and 'DHA' logo on the other side.

COMPOSITION:

Each Vasican tablet contains : Bromhexine HCl BP 8 mg/tablet
Excipients :
Corn Starch, Lactose Mesh, Acacia Powder, FDC Blue No. 1 Aluminum Lake HT, Quinoline Yellow E104, Magnesium Stearate.

PHARMACOLOGY :

VASICAN contains bromhexine HCl which is a mucolytic agent. It acts on the mucous gland in the respiratory tract to change the structure of bronchial secretions and to increase the volume and reduce the viscosity of the sputum.

PHARMACOKINETICS :

Oral administration Bromhexine HCl is absorbed rapidly from the gastrointestinal tract, with peak plasma concentration achieved in about one hour. It undergoes extensive first-pass metabolism in the liver and has an elimination half-life of 6.5 hours.

INDICATIONS :

For the reduction of sputum viscosity.

RECOMMENDED DOSAGE :

Adults and children over 12 years : 8 mg, 3 times a day
Children : 6 to 12 years : 4 mg, 3 times a day

WARNINGS AND PRECAUTIONS :

Bromhexine HCl should be given with caution to patient with gastric ulceration. In very few cases, severe skin disorders such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported after administration of mucolytic substances (e.g. bromhexine), which in most cases could be explained by the patient's underlying disease and/or the concomitant medication. In addition, during the early phase of such severe skin reactions, a patient may first experience non-specific influenza-like prodromes (e.g. fever, aching body, rhinitis, cough and sore throat). It is possible that a symptomatic treatment is started with a cough and cold preparation due to these misleading non-specific influenza-like prodromes. Therefore, if new skin or mucosal lesions occur, medical advice should be sought immediately and treatment with bromhexine discontinued as a precaution.

USE IN PREGNANCY AND LACTATION :

It is advisable to avoid use during the first three months of pregnancy and during lactation, Bromhexine HCl is excreted in breast milk.

DRUG INTERACTION :

Bromhexine HCl enhance the penetration of antibiotics (e.g. amoxycillin, cefuroxime, erythromycin and doxycycline) into the lung tissues. In elderly patients with acute exacerbation of chronic bronchopulmonary disease, a combination of cephalixin and bromhexine HCl shows that the symptoms

and ventilatory functions were improved to a greater extent in the group treated with antibiotic and mucolytic combination than with the antibiotic alone. No other clinically unfavourable interaction with medications have been reported.

CONTRAINDICATIONS :

Known hypersensitivity to bromhexine.

SIDE EFFECTS :

Gastrointestinal irritations, allergic reactions including skin rashes, bronchospasm, angio-oedema and anaphylaxis, transient increased serum amino transaminase.

Immune System Disorders

Frequency not known : Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorders

Frequency not known : Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

OVERDOSAGE AND TREATMENT :

To date, no symptoms of overdose have been reported.

STORAGE :

Protect from light.
Store below 30°C.

EXPIRY PERIOD :

Do not use after the expiry date stated on the label.

AVAILABILITY :

10x10x10's in blister pack

DATE OF REVISION : November 2021

For further information, please consult your physician or pharmacist.

Manufactured by:

PT Actavis Indonesia
Jl. Raya Bogor Km. 28, Kelurahan Pekayon, Kecamatan Pasar Rebo
Kota Jakarta Timur, Provinsi DKI Jakarta, Indonesia

Actavis
AAAN8699
41448-04

Product Name : Slip Vasican
Product Code : 41448-04
Dimension : 110 x 200 mm
Colour : ■ Black (1/1)
Paper : HVS 70 gr
Finishing : Lipat 3 x (11x5cm)
Date / Rev :