

BEACOLUX

TABLET 5mg ENTERIC COATED

DESCRIPTION

A 5.5mm in diameter, round, convex, enteric-coated yellow tablet.

INDICATION

Constipation, bowel evacuation before radiological procedures, endoscopy and surgery.

PHARMACODYNAMICS

Bisacodyl acts by increasing intestinal motility. It is thought to increase peristalsis by a direct effect on the smooth intestinal musculature by stimulation of intramural nerve plexi. Also have been shown to promote fluid and ion accumulation in the colon to increase the laxative effect.

PHARMACOKINETICS

Oral and rectal absorption of bisacodyl is minimal. Onset of action is about 1/4 to 1 hour. The absorbed doses is eliminated through renal.

DOSAGE

Adults: 2 tablets taken at bedtime in order to produce 1 to 2 actions the next morning. If taken half hour before breakfast, the effect will take place about 5 hours later. In stubborn cases of constipation the dose may be increased to 4 tablets.

Tablet should be swallowed whole, cannot be chewed or crush, should not be taken within 1 hour after taking an antacid or milk.

OR AS DIRECTED BY PHYSICIAN

CONTRAINDICATIONS

Intestinal obstruction that requires surgery.

PRECAUTIONS

Best avoided in pregnant women.

SIDE EFFECTS

Bisacodyl may cause abdominal discomfort such as colic or cramps. When administered rectally it sometimes causes irritation and repeated use may cause proctitis or sloughing of the epithelium. Prolonged use or overdosage can result in diarrhea with excessive loss of water and electrolytes. There is also the possibility of developing an atonic non-functioning colon.

PACKAGING

BEACOLUX tablet is packed in bottles of 30's. Each tablet contains 5mg bisacodyl.

STORAGE CONDITION

Store in a dry place below 30°C.

Keep away from light.

Keep out of reach of children.

Jauhi daripada kanak-kanak.

SHELF-LIFE

Please refer to box or label.

Route of administration: Oral

Product Registration Holder:

DUOPHARMA MANUFACTURING
(BANGI) SDN. BHD.

Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13,
Bangi Industrial Estate,
43650 Bandar Baru Bangi,
Selangor, Malaysia.

Manufacturer:

DUOPHARMA MANUFACTURING
(BANGI) SDN. BHD.

Lot No. 2 & 4, Jalan P/7, Section 13
Bangi Industrial Estate,
43650 Bandar Baru Bangi,
Selangor, Malaysia.



150000XXXX XX
Revision Date: April 2022

100%

128mm(h) x 128mm(w)