

ALUCID SUSPENSION
For Antacid and Anti-flatulent

INDICATIONS

- **Alucid Suspension** can be used for relief of gastric pain, hyperacidity, flatulence and heartburn.
- **Alucid Suspension** will relieve and control pain in the symptomatic management of gastric and duodenal ulcers and reflux oesophagitis by neutralising hydrochloric acid in gastric secretion.
- **Alucid Suspension** is also suitable as a domestic remedy in conditions affecting the stomach which may not necessarily be related to hyperacidity.

CONTENTS

Each 10ml (2 teaspoonful) of **Alucid Suspension** contains:-
- Dried Aluminium Hydroxide Gel 800mg (equiv to Aluminium Hydroxide 612mg)
- Magnesium Hydroxide 800mg
- Simethicone 60mg
Preservatives:-
Cetylpyridinium Chloride 10mg

DOSAGE

- Adult : 5 to 10ml (1 to 2 teaspoonful)
 - Children : 2.5 to 5 ml (1/2 to 1 teaspoonful)
- To be taken between meals and at bed time or as directed by physician.

PHARMACODYNAMICS

Aluminium Hydroxide, given by mouth, slowly reacts with the hydrochloric acid in the stomach to form soluble aluminium chloride, some of which is absorbed.

Magnesium Hydroxide is used in conjunction with aluminium containing antacids in order to reduce adverse gastro-intestinal effects and to produce a more sustain action.

Simethicone lowers surface tension and when administered by mouth causes bubbles of gas in the gastro-intestinal tract to coalesce, thus aiding their dispersion.

Alucid Suspension comes with a cooling and refreshing peppermint flavour.

Route of administration: Oral

PRODUCT DESCRIPTION

A white, mint flavoured homogeneous suspension.

SIDE EFFECTS

Constipation, diarrhoea, intestinal obstruction might happen if a larger than recommended dose of **Alucid Suspension** has been taken.

CONTRAINDICATION

Alucid Suspension should not be taken by persons who suffers from severe renal function impairment.

PRECAUTIONS

Use with care in renal insufficiency. Should not be used if symptoms of appendicitis are present. Do not take within 1 to 2 hours of other oral medication as it may impair their absorption. Please consult your physician if pain persists.

This product contains less than 1mmol sodium (23mg) per dose i.e. essentially 'sodium free'.

PACKING

100ml.

SHAKE WELL BEFORE USE

STORAGE

- **KEEP THE CONTAINER TIGHTLY CLOSED**
- **STORE IN A DRY PLACE BELOW 30°C**
- **PROTECT FROM LIGHT**
- **KEEP OUT OF REACH OF CHILDREN**

SHELF LIFE

Please refer to outer package

ALUCID 药水
缓和胃酸及气胀

说明

- **Alucid 药水**有助于减轻胃痛、胃酸过多、气胀和胃灼热。
- **Alucid 药水**有助于中和胃部的氮氯化酸，进而控制及减轻胃和十二指肠溃疡造成的疼痛以及食道逆流症状。
- **Alucid 药水**也适于治疗不是因胃酸过多引起的胃部疼痛／不适。

成份

每10毫升（2茶匙）**Alucid 药水**含有：
- 800 毫克 Dried Aluminium Hydroxide Gel
（相等予 612毫克 Aluminium Hydroxide）
- 800 毫克 Magnesium Hydroxide
- 60 毫克 Simethicone

防腐劑

Cetylpyridinium Chloride 10mg

劑量

- 成人：5至10毫升（1至2茶匙）
 - 儿童：2.5至5毫升（½至1茶匙）
- 在用餐之间及临睡前服用，或遵照医生指示。

副作用

若服用大量**Alucid 药水**，可能出现便秘、腹泻及肠胃阻塞现象。

禁忌

患有严重肾脏功能衰竭者请勿服用**Alucid 药水**。

警告

肾脏机能不全者应谨慎服用。如果出现阑尾炎症状，请勿服用。请勿服用其他口服药物之后的1至2小时内服用**Alucid 药水**，因为此举可能会削弱其吸收力。
如果疼痛持续，请即询问您的医生。

包装

100ml
服用前先摇匀。

贮藏

- 确保瓶盖紧关。
- 置于干燥及摄氏30°C 以下。
- 远离光线。
- 置于孩童无法触及之处。

贮藏期

请参阅盒内中文说明

ALUCID SUSPENSION
Sediaan Antasid dan Anti-flatulen

INDIKASI

- **Alucid Suspension** digunakan untuk melegakan kesakitan akibat gastrik, asid berlebihan, flatulen serta pedih di bahagian hulu hati.
- **Alucid Suspension** digunakan untuk melegakan dan mengawal kesakitan akibat gastrik dan ulser di bahagian bawah dan atas usus perut dengan meneutralkan asid hidroklorik di bahagian berkenaan.
- **Alucid Suspension** juga sesuai digunakan sebagai rawatan kepada sakit perut yang tidak berkaitan dengan kandungan asid yang berlebihan.

KANDUNGAN

Setiap 10ml (2 sudu teh) **Alucid Suspension** mengandungi :-
- Dried Aluminium Hydroxide Gel 800mg (equiv to Aluminium Hydroxide 612mg)
- Magnesium Hydroxide 800mg
- Simethicone 60mg

Bahan Pengawet :-

Cetylpyridinium Chloride 10mg

SUKATAN

- Dewasa: 5 hingga 10ml (1 hingga 2 sudu teh)
 - Kanak-kanak : 2.5 hingga 5ml (1/2 hingga 1 sudu teh)
- Diambil di antara waktu makan dan sebelum tidur atau seperti yang diarahkan oleh pakar perubatan.

KESAN SAMPINGAN

Sembelit, cirit-birit dan jangkitan usus tersumbat mungkin berlaku jika dos **Alucid Suspension** diambil secara berlebihan.

PERHATIAN

Alucid Suspension tidak boleh digunakan oleh pesakit buah pinggang yang serius.

AMARAN

Gunakan secara berhati-hati sekiranya anda mengalami masalah buah pinggang. Jangan gunakan sekiranya anda mempunyai tanda-tanda penyakit appendik. Tidak boleh diambil 1 atau 2 jam selepas mengambil ubat-ubat lain kerana ia mungkin memberi kesan terhadap keberkesanananya.
Silalah rujuk kepada pakar perubatan sekiranya sakit berlarutan.

BUNGKUSAN

100ml.
GONCANG SEBELUM DIGUNAKAN.

PENYIMPANAN

- **PASTIKAN BEKAS DITUTUP RAPAT.**
- **SIMPAN DI TEMPAT KERING DI BAWAH SUHU 30°C.**
- **JAUHKAN DARI CAHAYA.**
- **JAUHI DARIPADA KANAK-KANAK**



JANGKA HAYAT
Silalah rujuk pada pek luar.

Revision date: Mar 2026
1500010523 N.2

Product Registration Holder:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.
Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13, Bangi Industrial Estate,
43650 Bandar Baru Bangi, Selangor, Malaysia.

Manufacturer:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.
Lot No. 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate,
43650 Bandar Baru Bangi, Selangor Malaysia.