

Title: Mifuko Ointment - PI

Date: 04 Jul 2022

MIFUKO OINTMENT
(Skin Anti-infectives)

Active Ingredients:

Salicylic Acid 5% w/w
Zinc Oxide 6% w/w

Presentation:

Tube of 20g.

Descriptions:

White smooth ointment.

Pharmacology:

Salicylic Acid has bacteriostatic and fungicidal actions. It is applied externally in the treatment of persistent eczema, psoriasis, ringworm and other parasitic skin conditions. It also has keratolytic actions useful for dry scaling skin conditions. Zinc Oxide when applied externally acts as an astringent for the skin and as a soothing and protective application in eczema.

Indications:

For ringworm, athlete's foot, eczema, psoriasis, seborrhoeic dermatitis, scabies, warts and corns.

Directions of Use:

Clean and dry the affected areas. Apply 2 times daily.

Symptoms and Treatments of Overdose:

Symptoms: Symptoms of systemic salicylate poisoning varies from dizziness, tinnitus, deafness, sweating, headache, nausea and vomiting to hyperventilation, restlessness, ketosis, respiratory alkalosis, metabolic acidosis and depression of the central nervous system.

Treatments:

- Empty stomach by inducing emesis or by aspiration and lavage.
- Forced alkaline diuresis may be required and should be continued until the plasma salicylate concentration is less than 350 mcg per ml in adults, the intravenous fluids can be stopped and patient encouraged to take fluid by mouth.
- Plasma electrolytes especially potassium and the acid-base balance should be monitored regularly. Acidaemia must be corrected by infusion of sodium bicarbonate before starting forced diuresis.
- In the presence of cardiac or renal impairment or in very severe intoxication haemodialysis or peritoneal dialysis may be necessary.

Precautions:

For external use only. Avoid contact with eyes. Not to be taken.

Use with care on large areas of skin. Camphor can cause convulsions. It is contraindicated in infants below 2 years of age. Cautions must be exercised when older children are treated.

It is dangerous to place any camphor containing product into the nostrils of children. A small amount applied in this way may cause immediate collapse.

Undesirable Effects:

Skin dermatitis may occur on sensitive skin.

Storage:

Store below 30°C. Protect from light. Keep container tightly closed.

Keep out of reach of children.

Jauhkan daripada capaian kanak-kanak.

Shelf Life:

5 years from the date of manufacture.

Manufacturer and Product Registration Holder

WINWA MEDICAL SDN. BHD.

1952, Lorong I.K.S. Bukit Minyak 3,

Taman I.K.S. Bukit Minyak,

14000 Bukit Mertajam,

Pulau Pinang, Malaysia.

MAL19912829X BRU16032020NP

Date of Revision: 4 July 2022

X.EMK-1.09

Size: 55mm x 100mm
■ Dark Blue (Pantone 2767U)