

Important information. Please read carefully.

HYDROSIL

TABLET



Composition

Each chewable tablet contains Magnesium Hydroxide 200mg, Aluminium Hydroxide 200mg and Simethicone 20mg.

Product Description

White colour, round, flat bevelled edge tablet with single breakline on one side and crown logo on reverse, 12.5mm in diameter.

Pharmacodynamics

The antacid effect takes place in the stomach where both magnesium hydroxide and aluminium hydroxide, at the appropriate dose neutralize gastric acid production.

The constipating effect of aluminium hydroxide is counteracted by the laxative effect of magnesium hydroxide.

Simethicone has an antifatulent effect.

Pharmacokinetics

When administered orally, aluminium hydroxide and magnesium hydroxide react with the hydrochloric acid produced in the stomach to form salts, which are only partially absorbed.

Absorbed salts are excreted mainly in the urine.

Indications

Hydrosil is used for the relief of pain in gastric ulcer, gastric pain or discomfort due to excessive acid secretion in the stomach (dyspepsia) and in reflux oesophagitis (heartburn).

Dosage and Administration

2 - 4 tablets to be sucked or chewed before swallowing when symptoms occur or likely to occur, usually between meals and at bedtime.

Contraindications

Hydrosil is contraindicated in patients with gastrointestinal obstruction, renal insufficiency and in patients with low plasma concentration of phosphate (hypophosphataemia).

Warnings and Precautions

Avoid prolonged use. Hydrosil should be used with caution in patients on dialysis and patients with low-phosphate diets.

Interactions with other medicaments

Like other antacids, Hydrosil should not be taken together with other drugs such as tetracycline, iron, rifampicin, digoxin, cimetidine, chlorpromazine, penicillamine; as it may reduce the effects of these drugs. If prescribed, they may be taken one to two hours after Hydrosil.

Pregnancy and Lactation

There are no available data for use in pregnant women. This product should only be used if absolutely necessary, under direct medical supervision.

When taken in accordance with the indicated dosage regimen, aluminum hydroxide and its magnesium salt combinations are considered compatible with breast-feeding because of limited maternal absorption.

Side Effects

At the recommended dosage, Hydrosil is usually well tolerated.

Immune system disorders:

Frequency not known: angioedema, anaphylactic reactions, hypersensitivity reactions, urticaria, pruritus.

Gastrointestinal disorders:

Uncommon (≥1/1 000 to <1/100): diarrhoea or constipation.

Frequency not known: abdominal pain.

Metabolism and nutrition disorders:

Very rare (<1/10 000): hypermagnesaemia (with prolonged use in patients with kidney impairment).

Frequency not known: Hyperaluminemia, hypophosphatemia in prolonged use or at high doses, or even normal doses in patients on low-phosphorus diets, or in infants, which may result in increased bone resorption, hypercalciuria or osteomalacia.

Symptoms and Treatment of Overdose

At the recommended dosage, Hydrosil is usually well tolerated. In the event of severe overdosage or poisoning, hypermagnesaemia may occur. Hospital admission is strongly advised for patients showing symptoms of overdosage; which may include flushing of the skin, thirst, hypotension, drowsiness, respiratory depression, cardiac arrhythmia, coma and cardiac arrest.

Hypermagnesaemia may be corrected by intravenous injection of 10 to 20ml of 10% Calcium Gluconate. Dialysis and correction of fluid deficit may be of value.

Storage Condition

Store below 30°C.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN.

JAUHI DARI KANAK-KANAK.

Shelf Life

The shelf life is indicated on the packaging.

Pack Size

Available in blister pack of 10 x 10's and 100 x 10's.

Not all pack size(s) is available for sale.

For further information, please consult your pharmacist or physician.

Revision Date: 15-Oct-2025

Manufacturer and Product Registration Holder (in Malaysia):

Xepa-Soul Pattinson (Malaysia) Sdn. Bhd.

1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.