

Important Information. Please read carefully.

Bacidin™ 5%

Concentrate Solution

Composition

Contains Chlorhexidine Gluconate 5 % w/v

Pharmacodynamics

Chlorhexidine is a disinfectant which is effective against a wide range of vegetative Gram-positive and Gram-negative bacteria.

Pharmacokinetics

Chlorhexidine is poorly absorbed through skin.

Indication

Bacidin is used for the pre-operative disinfection of the skin, as a disinfectant for wound and burns, obstetrics and gynaecology. It is also used for storage of sterile instruments and for sterilisation of aqueous solutions.

Dosage

FOR EXTERNAL USE ONLY.

For disinfection of wounds and burns; swabbing in obstetrics & gynaecology, and storage of sterile instruments: 0.05 % w/v aqueous solution.

For pre-operative skin disinfection, emergency disinfection by 2 minutes immersion (excluding endoscope) : 0.5 % w/v Alcoholic solution.

Sterilisation of aqueous solutions : 1 % or less of **Bacidin**. Autoclave at 115°C for 30 minutes or 121°C for 15 minutes.

Dilution guide:

0.05 % w/v Aqueous solution : Dilute 10 ml of **Bacidin** with water up to 1L

0.5 % w/v Alcoholic solution : Dilute 100 ml of **Bacidin** with 200 ml of water and made up to 1 L with 95% alcohol.

Solutions for storage of instruments over 8 hours should contain 0.1% of sodium nitrite to prevent corrosion. Change solutions fortnightly.

Contraindications

Bacidin should not be used on the brain, meninges, or perforated ear-drum. Syringes and needles that have been immersed in **Bacidin** solutions should be thoroughly rinsed with Water for Injections before use for spinal injection.

Warnings and Precautions

Do not use undiluted solutions on sensitive tissues.

Avoid contact with the eyes, brain, meninges and middle ear. Chlorhexidine is incompatible with cork, soaps and other anionic materials.

Do not mix with other chemicals or detergents.

Allergy Alert

Bacidin contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare. **Bacidin** should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine. If any signs or symptoms of suspected hypersensitivity reaction such as itching, skin rash, hives, redness, swelling (including facial swelling), breathing difficulties/ wheezing, shock, light headedness, and rapid heart rate develop, immediately stop using the product and seek medical help. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Side Effects

Occasionally, skin sensitivity may occur.

Immune system disorders

Frequency not known : Hypersensitivity including anaphylactic shock.

Symptoms and Treatment for Overdosage

Accidental ingestion may cause irritation of gastrointestinal tract. Chlorhexidine is poorly absorbed from the gastrointestinal tract. Toxic symptoms may include dyspnoea and cyanosis due to paralysis of the respiratory muscles.

First Aids

Treatment may include the immediate administration of milk and gastric lavage given one to two hours later. Supportive and symptomatic treatment may be given.

Shelf-life

The expiry date is indicated on the packaging.

Presentation

Clear, red solution in packs of 500 ml.

Storage

Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN/ JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date : 01-Dec-2017



Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.