

PEDIVITE FORTE CAPSULES

Composition:

Each soft gelatin capsule contains:

Substance Name	Quantity in mg Per Capsule	% USRDA
Vitamin A U.S.P. (As Palmitate)	2000IU	40%
Vitamin B-1 B. P.	1 mg	66%
Vitamin B-2 B. P.	1 mg	58%
Vitamin B-6 B. P.	1.34 mg	67%
Niacinamide B. P.	8 mg	40%
Calcium Pantothenate U.S.P.	1 mg	10%
Vitamin B-12 B. P.	1 mcg	16%
Ascorbic Acid B. P.	33.3 mg	55.5%
Vitamin D-3 U.S.P.	270 IU	67.5%
Calcium (From Dicalcium Phosphate B. P.)	190 mg	19%
Phosphorous (Dicalcium Phosphate B. P.)	140 mg	27%
Iron (From Ferrous Sulphate B. P.)	5 mg	25%
Copper (From Copper Sulphate B. P.)	0.01 mg	0.5%
Iodine (From Potassium Iodide B. P.)	0.0015 mg	0.05%
Magnesium (From Magnesium Sulphate B. P.)	1.0 mg	0.25 %
Manganese (From Manganese Sulphate B.P.)	0.01 mg	0.25%
Potassium (From Potassium Iodide B. P.)	1 mg	-
Zinc (From Zinc Sulphate B. P.)	0.15 mg	1%
Preservatives:		--
Butylated Hydroxytoluene	0.100mg	
Sodium Methyl paraben	1.00 mg	--
Sodium propyl paraben	0.100mg	--
Excipients	q.s.	

Source of capsule: Bovine.

Product Description:

Brown coloured oblong shaped soft gelatin capsules.

Indications:

Pedivite forte capsules is indicated for food dietary supplement,

Dose/Use Instruction:

One capsule daily with water

Contra – Indication:

Haemochromatosis and Haemosiderosis are contra-indications to iron therapy.

Warning / Precautions:

Certain body conditions may require additional-nutritional supplementation, during pregnancy, for instance supplementation with fat soluble vitamins and minerals may be required according to the dietary habits of the individual. Pedivite forte is not intended for treatment of severe specific deficiencies.

Drug Interaction:

There is no report on drug interactions reported.

Side Effects:

Excess overdosage would cause condition known as hypervitaminosis may occur.

Signs and Symptoms of Overdose and Treatment:

Amounts higher than recommended dosage have been found to be toxic orally. Toxicity manifestations depend on the age, dosage size and duration of administration. A condition known as hypervitaminosis may occur may be controlled on discontinuation.

Storage Condition

Store below 25° C. Protect from light and moisture

Shelf Life

3 years from date of Manufacture.

Presentation:

Packs of 500's capsules in a plastic container;

Packs of 30's capsules in a plastic container;

Packs of blister of 7 capsules, 3 Blisters are packed in a carton;

Manufactured by:

Unique Pharm Labs,

83 B & C Dr. Annie Besant Rd,

Bombay 400 018, India.