



ARTWORK

Please return to:
LEO Pharma A/S
Internal Market Access
Industriparken 55 – DK-2750 – Ballerup

Scale 100%	Get-up HK/LK/MY/SG	Item No. 000000	Rev. No. XX	Sent by e-mail ▼	
Subject INS 145 x 148 mm			Date 04/10/13	Date	Date
Colour Black			Sign. JUG	Sign.	Sign.

Preparation Strength Fucidin® cream	Supplier / Place of production Ireland
---	---

Comments:

Page 1 of 2 Pharmacode XX Pointsize: 9,5 pt Mock-up for reg. purpose

145 mm

148 mm

Fucidin®

Plain topical antibiotic

Cream
Fusidic acid 20 mg/g in neutral cream base.

Pack Sizes: 5g, 15g.

Properties
Fusidic acid is an antibiotic with powerful antibacterial activity against a number of gram-positive organisms. Staphylococci, including the strains resistant to penicillin or to other antibiotics, are particularly susceptible to fusidic acid. Besides pronounced antibacterial activity against the majority of organisms responsible for skin infections, Fucidin® cream also possesses unique ability for penetration to the focus of infection, even when applied to intact skin.

Indications
Fucidin® cream is indicated for treatment of skin infections caused by staphylococci, streptococci, Propionibacterium acnes, Corynebacterium minutissimum, and other organisms sensitive to fusidic acid. The most important indications being:

Impetigo	Boils	Carbuncles	Paronychia
Infected wounds	Sycosis barbae	Hidradenitis	Erythrasma
Folliculitis			

Fucidin® cream is an invisible, non-staining preparation, which is cosmetically acceptable for treatment of face and scalp infections.

Application
Fucidin® cream is applied to the affected area 2–3 times daily, generally for a period of 7 days, except in treatment of acne where therapy should be extended for a longer period according to the nature of this disease.

XX-000000

000000-00

Item-numb
See details

80 mm

Space for text: 125 x 128 mm.

2. PROOF FROM	JUG	Mock-up Approval Stamp (MAS)		
Date	04/10/13	Graphic Design	Editorial Proof	Second Approver
New proof requested ☐	Sign.:	According to: Instruction For Use, Mock-up Approval Stamp (MAS) SOP_000647 and SOP_000962 ☐	According to: Instruction For Use, Mock-up Approval Stamp (MAS) SOP_000647 and SOP_000962 ☐	Product name ☐
		Sign.: Date:	Sign.: Date:	Dosage form ☐
Date:				Strength/Stripes ☐
				Pack size ☐
				Prompts ☐
				Item No./Reg. No. ☐
				Sign.: Date:



ARTWORK

Please return to:
LEO Pharma A/S
Internal Market Access
Industriparken 55 – DK-2750 – Ballerup

Scale	Get-up	Item No.	Rev. No.	Sent by e-mail	
100%	HK/LK/MY/SG	000000	XX	▼	
Subject	INS 145 x 148 mm		Date	Date	Date
			04/10/13		
Colour	Black		Sign.	Sign.	Sign.
			JUG		

Preparation Strength	Fucidin® cream	Supplier / Place of production	Ireland
-------------------------	----------------	--------------------------------	---------

Comments:
Page 2 of 2

145 mm		
148 mm	Contraindications Infections caused by non-susceptible organisms. Patients with hypersensitivity to any of the ingredients. Pregnancy and lactation There is inadequate evidence of safety in human pregnancy. Animal studies and many years of clinical experience have suggested that fusidic acid is devoid of teratogenic effects. Safety in nursing mothers has not been established. Side-effects Fucidin® cream is remarkably well tolerated, and there is an extremely low frequency of hypersensitivity reactions. Drug interaction Not relevant. Incompatibilities Fucidin® cream is a ready-for-use preparation and should, therefore, not be mixed with other ingredients or preparations. Storage Do not store above 30°C. Manufactured by: LEO Laboratories Ltd 285, Cashel Road, Crumlin, Dublin 12, Ireland This leaflet was last revised in August 2013. LEO Pharma A/S – Ballerup – Denmark LEO	80 mm XX-000000 000000-00 Item-numb See details

Space for text: 125 x 128 mm.